

Notes on homotopical algebra

Z.L. Low

2nd March 2013

PREFACE

These notes are intended as a kind of annotated index to the various standard references in homotopical algebra: the focus is on definitions and statements of results, *not* proofs.

CONTENTS

Foundations	I
0.1. Set theory	I
0.2. Accessibility and ind-completions	10
0.3. Change of universe	23
Simplicial sets	33
1.1. Basics	33
1.2. The Quillen model structure	39
Homotopical categories	41
2.1. Basics	41
2.2. Homotopical Kan extensions	42
Model categories	45
3.1. Basics	45
3.2. Left and right homotopy	49
3.3. The homotopy category	54

CONTENTS

Generalities	57
A.1. Factorisation systems	57
A.2. Relative categories	64
A.3. Kan extensions	70
Bibliography	71
Index	75

FOUNDATIONS

0.1 Set theory

In category theory it is often convenient to invoke a certain set-theoretic device commonly known as a ‘Grothendieck universe’, but we shall say simply ‘universe’, so as to simplify exposition and proofs by eliminating various circumlocutions involving cardinal bounds, proper classes etc.

Definition 0.1.1. A **pre-universe** is a set \mathbf{U} satisfying these axioms:

1. If $x \in y$ and $y \in \mathbf{U}$, then $x \in \mathbf{U}$.
2. If $x \in \mathbf{U}$ and $y \in \mathbf{U}$ (but not necessarily distinct), then $\{x, y\} \in \mathbf{U}$.
3. If $x \in \mathbf{U}$, then $\mathcal{P}(x) \in \mathbf{U}$, where $\mathcal{P}(x)$ denotes the set of all subsets of x .
4. If $x \in \mathbf{U}$ and $f : x \rightarrow \mathbf{U}$ is a map, then $\bigcup_{i \in x} f(i) \in \mathbf{U}$.

A **universe** is a pre-universe \mathbf{U} with this additional property:

5. $\omega \in \mathbf{U}$, where ω is the set of all finite (von Neumann) ordinals.

Example 0.1.2. The empty set is a pre-universe, and with very mild assumptions, so is the set \mathbf{HF} of all hereditarily finite sets.

¶ **0.1.3.** The notion of universe makes sense in any material set theory, but their existence must be postulated. We adopt the following:

- **Grothendieck–Verdier universe axiom.** For each set x , there exists a universe \mathbf{U} with $x \in \mathbf{U}$.

For definiteness, we may take our base theory to be Mac Lane set theory, which is a weak subsystem of Zermelo–Fraenkel set theory with choice (ZFC). Readers interested in the details of Mac Lane set theory are referred to [Mathias, 2001], but in practice as long as one is working at all times *inside some universe*, one may as well be working in ZFC. Indeed:

Proposition 0.1.4. *With the assumptions of Mac Lane set theory, any universe is a transitive model of ZFC.*

Proof. Let \mathbf{U} be a universe. By definition, \mathbf{U} is a transitive set containing pairs, power sets, unions, and ω , so the axioms of extensionality, empty set, pairs, power sets, unions, choice, and infinity are all automatically satisfied. We must show that the axiom schemas of separation and replacement are also satisfied, and in fact it is enough to check that replacement is valid; but this is straightforward using axioms 2 and 4. ■

Definition 0.1.5. Let \mathbf{U} be a pre-universe. A **U-set** is a member of \mathbf{U} , a **U-class** is a subset of \mathbf{U} , and a **proper U-class** is a **U-class** that is not a **U-set**.

Lemma 0.1.6. *A U-class X is a U-set if and only if there exists a U-class Y such that $X \in Y$.* ■

Proposition 0.1.7. *If \mathbf{U} is a universe, then the collection of U-classes is a transitive model of Morse–Kelley class–set theory (MK), and so is a transitive model of von Neumann–Bernays–Gödel class–set theory (NBG) in particular.* ■

Definition 0.1.8. A **U-small category** is a category \mathbb{C} such that $\text{ob } \mathbb{C}$ and $\text{mor } \mathbb{C}$ are **U-sets**. A **locally U-small category** is a category \mathcal{D} satisfying these conditions:

- $\text{ob } \mathcal{D}$ and $\text{mor } \mathcal{D}$ are **U-classes**, and
- for all objects x and y in \mathcal{D} , the hom-set $\mathcal{D}(x, y)$ is a **U-set**.

An **essentially U-small category** is a category \mathcal{D} for which there exist a **U-small category** \mathbb{C} and a functor $\mathbb{C} \rightarrow \mathcal{D}$ that is fully faithful and essentially surjective on objects.

Proposition 0.1.9. *If \mathbb{C} is a U-small category and \mathcal{D} is a locally U-small category, then the functor category $[\mathbb{C}, \mathcal{D}]$ is locally U-small.*

Proof. Strictly speaking, this depends on the set-theoretic implementation of ordered pairs, categories, functors, etc., but at the very least $[\mathbb{C}, \mathbb{D}]$ should be isomorphic to a locally \mathbf{U} -small category.

In the context of $[\mathbb{C}, \mathbb{D}]$, we may regard functors $\mathbb{C} \rightarrow \mathbb{D}$ as being the pair consisting of the *graph* of the object map $\text{ob } \mathbb{C} \rightarrow \text{ob } \mathbb{D}$ and the *graph* of the morphism map $\text{mor } \mathbb{C} \rightarrow \text{mor } \mathbb{D}$, and these are \mathbf{U} -sets by the \mathbf{U} -replacement axiom. Similarly, if F and G are objects in $[\mathbb{C}, \mathbb{D}]$, then we may regard a natural transformation $\alpha : F \Rightarrow G$ as being the triple (F, G, A) , where A is the set of all pairs (c, α_c) . ■

One complication introduced by having multiple universes concerns the existence of (co)limits.

Theorem 0.I.10 (Freyd). *Let \mathcal{C} be a category and let κ be a cardinal such that $|\text{mor } \mathcal{C}| \leq \kappa$. If \mathcal{C} has products for families of size κ , then any two parallel morphisms in \mathcal{C} must be equal.*

Proof. Suppose, for a contradiction, that $f, g : X \rightarrow Y$ are distinct morphisms in \mathcal{C} . Let Z be the product of κ -many copies of Y in \mathcal{C} . The universal property of products implies there are at least 2^κ -many distinct morphisms $X \rightarrow Z$; but $\mathcal{C}(X, Z) \subseteq \text{mor } \mathcal{C}$, so this is an absurdity. ■

Definition 0.I.11. Let \mathbf{U} be a pre-universe. A **\mathbf{U} -complete** (resp. **\mathbf{U} -cocomplete**) **category** is a category \mathcal{C} with the following property:

- For all \mathbf{U} -small categories \mathbb{D} and all diagrams $A : \mathbb{D} \rightarrow \mathcal{C}$, a limit (resp. colimit) of A exists in \mathcal{C} .

We may instead say \mathcal{C} has all **finite limits** (resp. **finite colimits**) in the special case $\mathbf{U} = \mathbf{HF}$.

Proposition 0.I.12. *Let \mathcal{C} be a category and let \mathbf{U} be a non-empty pre-universe. The following are equivalent:*

- (i) \mathcal{C} is \mathbf{U} -complete.
- (ii) \mathcal{C} has all finite limits and products for all families of objects indexed by a \mathbf{U} -set.

(iii) For each \mathbf{U} -small category \mathbb{D} , there exists an adjunction

$$\Delta \dashv \lim_{\longleftarrow \mathbb{D}} : [\mathbb{D}, C] \rightarrow C$$

where ΔX is the constant functor with value X .

Dually, the following are equivalent:

(i') C is \mathbf{U} -cocomplete.

(ii') C has all finite colimits and coproducts for all families of objects indexed by a \mathbf{U} -set.

(iii') For each \mathbf{U} -small category \mathbb{D} , there exists an adjunction

$$\lim_{\longrightarrow \mathbb{D}} \dashv \Delta : C \rightarrow [\mathbb{D}, C]$$

where ΔX is the constant functor with value X .

Proof. This is a standard result; but we remark that we do require a sufficiently powerful form of the axiom of choice to pass from (ii) to (iii). \square

¶ 0.I.13. In the **explicit universe convention**, the words ‘set’, ‘class’, etc. have their usual meanings, and in the **implicit universe convention**, these instead abbreviate ‘ \mathbf{U} -set’, ‘ \mathbf{U} -class’, etc. for a fixed (but arbitrary) universe \mathbf{U} . However, the word ‘category’ always refers to a category that is contained in *some* universe, which may or may not be \mathbf{U} . In all subsequent chapters, the implicit universe convention should be assumed *unless otherwise stated*.

We now recall some definitions and results about ordinal and cardinal numbers. Readers familiar with axiomatic set theory may wish to skip ahead.

Definition 0.I.14. A **von Neumann ordinal** is a set α with the following properties:

- If $x \in y$ and $y \in \alpha$, then $x \in \alpha$.
- The binary relation \in is strict total ordering of α .
- If S is a subset of α such that

$$- \emptyset \in S,$$

- If $\beta \in S$ and $\beta \cup \{\beta\} \in \alpha$, then $\beta \cup \{\beta\} \in S$.
- If $T \subseteq S$, then $\bigcup T \in S$.

then $S = \alpha$.

We identify 0 with the von Neumann ordinal \emptyset , and by induction, we identify the natural number $n + 1$ with the von Neumann ordinal $\{0, \dots, n\}$.

Proposition 0.I.15.

- (i) *If α is a von Neumann ordinal, then every member of α is an initial segment of α and is in particular a von Neumann ordinal.*
- (ii) *If α is a von Neumann ordinal, so is $\alpha \cup \{\alpha\}$. (This is usually denoted by $\alpha + 1$ and called the **successor** of α .)*
- (iii) *The union of a set S of von Neumann ordinals is another von Neumann ordinal. (This is usually denoted by $\sup S$ and called the **supremum** of S .)*
- (iv) *If \mathbf{U} is a pre-universe and $\kappa(\mathbf{U})$ is the set of von Neumann ordinals in \mathbf{U} , then $\kappa(\mathbf{U})$ a von Neumann ordinal, but $\kappa(\mathbf{U}) \notin \mathbf{U}$.*

Proof. Claims (i) – (iii) are all easy, and claim (iv) is Burali-Forti’s paradox. \square

Theorem 0.I.16 (Classification of well-orderings).

- (i) *In Zermelo–Fraenkel set theory, every well-ordered set is isomorphic to a unique von Neumann ordinal.*
- (ii) *In Mac Lane set theory, if \mathbf{U} is a pre-universe and X is a well-ordered set in \mathbf{U} , then X is isomorphic to a unique von Neumann ordinal in \mathbf{U} .*

Proof. Claim (i) is a standard result in axiomatic set theory, and claim (ii) is an obvious corollary. \square

Definition 0.I.17. A **transitive set** is a set T such that, given $x \in y$, if $y \in T$, then $x \in T$ as well. The **transitive closure** of a set X is a set $\text{tcl}(X)$ such that, for all transitive sets T with $X \subseteq T$, we have $\text{tcl}(X) \subseteq T$ as well.

Lemma 0.I.18. *In Mac Lane set theory, every set has a unique transitive closure.*

Proof. One of the axioms of Mac Lane set theory states that every set X is a member of some transitive set T , and so $X \subseteq T$. Clearly, the intersection of any family of transitive sets containing X is again a transitive set containing X , so $\text{tcl}(X)$ exists and is unique so long as there is at least one transitive set containing X . ■

Definition 0.1.19. A **partial rank function** from a transitive set T to a well-ordered set W is a partial function $\rho : T \rightarrow W$ with these properties:

- If $\emptyset \in T$, then $\rho(\emptyset)$ is the least element of W .
- If $y \in T$ and $\rho(x)$ is defined for all $x \in y$, then

$$\rho(y) = \min \{w \in W \mid \forall x \in y. \rho(x) < w\}$$

provided the RHS is defined.

- Otherwise $\rho(y)$ is undefined.

A **total rank function** is a partial rank function that is defined on its entire domain. The **rank** of a set X , if it exists, the least von Neumann ordinal $\text{rank}(X)$ for which there exists a total rank function $\text{tcl}(X) \rightarrow \text{rank}(X)$.

Proposition 0.1.20. *In Mac Lane set theory:*

- (i) *If T is a transitive set and W is a well-ordered set, then there is a unique partial rank function $\rho : T \rightarrow W$.*
- (ii) *If \mathbf{U} is a pre-universe and $x \in \mathbf{U}$, then $\text{rank}(x)$ can be defined by a Δ_0 -formula with \mathbf{U} as a parameter; and for each von Neumann ordinal α in \mathbf{U} , the set*

$$\mathbf{V}_\alpha = \{x \in \mathbf{U} \mid \text{rank}(x) < \alpha\}$$

is a \mathbf{U} -set.

- (iii) *Assuming the Grothendieck–Verdier universe axiom, $\text{rank}(x)$ is defined for all x .*

Proof. (i). This is a straightforward application of well-founded induction.

(ii). \mathbf{U} is a transitive set and the set $\kappa(\mathbf{U})$ of all von Neumann ordinals in \mathbf{U} is well-ordered by inclusion, so by claim (i) there is a partial rank function $\rho :$

$\mathbf{U} \rightarrow \kappa(\mathbf{U})$. ZFC proves that every set has a rank, so ρ must in fact be a total rank function; hence, for any $x \in \mathbf{U}$, $\text{rank}(x)$ is defined. It is clear that ρ can be defined by a Δ_0 -formula with only \mathbf{U} as a parameter, and the rest of the claim follows.

(iii). Obvious, assuming claim (ii). □

Definition 0.I.21. Two sets are **equinumerous** if there exists a bijection between them. A **cardinality class** in a pre-universe \mathbf{U} is an equivalence class under the relation of equinumerosity.

Definition 0.I.22. An **\aleph -number** is an infinite von Neumann ordinal κ such that, for any von Neumann ordinal λ such that κ and λ are equinumerous, we have $\kappa \subseteq \lambda$.

Example 0.I.23. The first infinite von Neumann ordinal, i.e. $\omega = \{0, 1, 2, \dots\}$, is the \aleph -number \aleph_0 .

Theorem 0.I.24 (Classification of cardinalities).

- (i) *In Zermelo–Fraenkel set theory, for every well-ordered infinite set X , there exists a unique \aleph -number κ such that X and κ are equinumerous.*
- (ii) *In Zermelo–Fraenkel set theory with the axiom of choice, the same is true for any infinite set whatsoever.*
- (iii) *In Mac Lane set theory, if \mathbf{U} is a universe and X is an infinite set in \mathbf{U} , then there exists a unique \aleph -number κ in the cardinality class of X .*
- (iv) *In Mac Lane set theory with the Grothendieck–Verdier universe axiom, if \mathbf{U} is a pre-universe and κ is an \aleph -number not in \mathbf{U} , then the cardinality of \mathbf{U} is at most κ .*

Proof. Claim (i) is a standard fact, whence claims (ii) and (iii), by the well-ordering theorem. Claim (iv) can be proven using axiom 4 for pre-universes. □

¶ **0.I.25.** Henceforth, we identify the cardinality class of a finite set with the unique von Neumann ordinal contained in that class, and similarly we identify the cardinality class of an infinite set with the unique \aleph -number in that class. These are the **cardinal numbers**.

Definition 0.1.26. A **cofinal subset** of a partially-ordered set X is a subset $Y \subseteq X$ such that, for all x in X , there exists some y in Y such that $x \leq y$. A **regular cardinal number** is an \aleph -number κ such that any cofinal subset of κ has cardinality equal to κ . A **singular cardinal number** is an \aleph -number that is not regular.

The following helps to motivate the definition of regular cardinal numbers.

Definition 0.1.27. Let \mathbf{U} be a pre-universe. An **arity class** in \mathbf{U} is a \mathbf{U} -class K of cardinal numbers satisfying the following conditions:

- $1 \in K$.
- If $\kappa \in K$ and $\lambda : \kappa \rightarrow K$ is a function, then the cardinal sum $\sum_{\alpha \in \kappa} \lambda(\alpha)$ is also in K .
- If $\kappa \in K$ and $\lambda : \kappa \rightarrow \mathbf{U}$ is a function such that each $\lambda(\alpha)$ is a cardinal number and $\sum_{\alpha \in \kappa} \lambda(\alpha) \in K$, then $\lambda(\alpha) \in K$ as well.

Theorem 0.1.28 (Classification of arity classes). *In Mac Lane set theory, if K is an arity class in a pre-universe \mathbf{U} , then K must be either*

- $\{1\}$, or
- $\{0, 1\}$, or
- of the form $\{\lambda \in \mathbf{U} \mid \lambda \text{ is a cardinal number and } \lambda < \kappa\}$ for some regular cardinal number κ (possibly not in \mathbf{U}).

Proof. The notion of arity class and this result are due to Shulman [2012]. \square

Definition 0.1.29. Let κ be a regular cardinal number. A κ -**small category** is a category \mathbf{C} such that $\text{mor } \mathbf{C}$ has cardinality *less than* κ . A **finite category** is an \aleph_0 -small category, i.e. a category \mathbf{C} such that $\text{mor } \mathbf{C}$ is finite. A **finite diagram** (resp. κ -**small diagram**, **U-small diagram**) in a category \mathbf{C} is a functor $\mathbb{D} \rightarrow \mathbf{C}$ where \mathbb{D} is a finite (resp. κ -small, \mathbf{U} -small) category.

Theorem 0.1.30. *Let \mathbf{U} be a pre-universe, and let \mathbf{U}^+ be a universe with $\mathbf{U} \in \mathbf{U}^+$. Let \mathbf{Set} be the category of \mathbf{U} -sets, and let \mathbf{Set}^+ be the category of \mathbf{U}^+ -sets.*

- (i) *If $X : \mathbb{D} \rightarrow \mathbf{Set}$ is a \mathbf{U} -small diagram, then there exist a limit and a colimit for X in \mathbf{Set} .*

- (ii) *The inclusion $\mathbf{Set} \hookrightarrow \mathbf{Set}^+$ is fully faithful and preserves limits and colimits for all \mathbf{U} -small diagrams.*

Proof. One can construct products, equalisers, coproducts, coequalisers, and hom-sets in a completely explicit way, making the preservation properties obvious. □

Corollary 0.1.31. *The inclusion $\mathbf{Set} \hookrightarrow \mathbf{Set}^+$ reflects limits and colimits for all \mathbf{U} -small diagrams.* ■

Corollary 0.1.32. *For any \mathbf{U} -small category \mathbb{C} :*

- (i) *The functor category $[\mathbb{C}, \mathbf{Set}]$ is \mathbf{U} -complete and \mathbf{U} -cocomplete, with limits and colimits for \mathbf{U} -small diagrams computed componentwise in \mathbf{Set} .*
- (ii) *The inclusion $[\mathbb{C}, \mathbf{Set}] \hookrightarrow [\mathbb{C}, \mathbf{Set}^+]$ is fully faithful and both preserves and reflects limits and colimits for all \mathbf{U} -small diagrams.* ■

Definition 0.1.33. An **strongly inaccessible cardinal number** is a regular cardinal number κ such that, for all sets X of cardinality less than κ , the power set $\mathcal{P}(X)$ is also of cardinality less than κ .

Example 0.1.34. \aleph_0 is a strongly inaccessible cardinal number and is the only one that can be proven to exist in ZFC. It is more conventional to exclude \aleph_0 from the definition of strongly inaccessible cardinal number by demanding that they be uncountable.

Proposition 0.1.35. *In Mac Lane set theory:*

- (i) *If \mathbf{U} is a non-empty pre-universe, then there exists a strongly inaccessible cardinal number κ such that the members of \mathbf{U} are all the sets of rank less than κ . Moreover, this κ is the rank and the cardinality of \mathbf{U} .*
- (ii) *If \mathbf{U} is a universe and κ is a strongly inaccessible cardinal number such that $\kappa \in \mathbf{U}$, then there exists a \mathbf{U} -set \mathbf{V}_κ whose members are all the sets of rank less than κ , and \mathbf{V}_κ is a pre-universe.*
- (iii) *If \mathbf{U} and \mathbf{U}' are pre-universes, then either $\mathbf{U} \subseteq \mathbf{U}'$ or $\mathbf{U}' \subseteq \mathbf{U}$; and if $\mathbf{U} \subsetneq \mathbf{U}'$, then $\mathbf{U} \in \mathbf{U}'$.*

Proof. (i). Let κ be the set of all von Neumann ordinals in \mathbf{U} ; this exists by Δ_0 -separation applied to \mathbf{U} . Since \mathbf{U} is closed under power sets and internally-indexed unions, κ must be a strongly inaccessible cardinal.

We can construct the set all of \mathbf{U} -sets of rank less than κ using transfinite recursion on κ as follows: starting with $\mathbf{V}_0 = \emptyset$, for each von Neumann ordinal α less than κ , we set $\mathbf{V}_{\alpha+1} = \mathcal{P}(\mathbf{V}_\alpha)$, and for each ordinal λ that is not a successor, we set $\mathbf{V}_\lambda = \bigcup_{\alpha < \lambda} \mathbf{V}_\alpha$. The well-foundedness of \in (restricted to \mathbf{U}) implies that in fact this must be all of \mathbf{U} .

Clearly, every set of rank less than κ is in fact a \mathbf{U} -set, and \mathbf{U} is itself a set of rank κ . The cardinality of \mathbf{U} is also κ , since κ is a regular cardinal number and any cardinal number less than κ is a member of \mathbf{U} .

(ii). We may construct \mathbf{V}_κ using the same method as in (i). By construction \mathbf{V}_κ satisfies axiom 1; since κ is infinite, \mathbf{V}_κ satisfies axioms 2 and 3; and since κ is strongly inaccessible, \mathbf{V}_κ satisfies axiom 4. Thus \mathbf{V}_κ is a pre-universe.

(iii). Again, let κ be the rank of \mathbf{U} . If $\kappa \in \mathbf{U}'$ then we can show by transfinite induction that $\mathbf{V}_\kappa \in \mathbf{U}'$ and so $\mathbf{U} \subsetneq \mathbf{U}'$; else we must have $\mathbf{U}' \subseteq \mathbf{V}_\kappa = \mathbf{U}$. ■

0.2 Accessibility and ind-completions

Prerequisites. § 0.1.

A classical technology for controlling size problems in category theory, due to Gabriel and Ulmer [1971], is the notion of accessibility. Though we make use of universes, accessibility remains important and is a crucial tool in verifying the stability of various universal constructions when one passes from one universe to a larger one.

Definition 0.2.1. Let κ be a regular cardinal. A κ -**filtered category** is a category \mathcal{J} satisfying these conditions:

- \mathcal{J} is **inhabited**, i.e. there exists an object in \mathcal{J} .
- If λ is a cardinal number strictly less than κ and S is a subset of $\text{ob } \mathcal{J}$ of cardinality λ , then there exist an object j and arrows $f_i : i \rightarrow j$ for each object i in S .
- If $f, g : i \rightarrow j$ are a pair of parallel arrows in \mathcal{J} , then there exist an object k and an arrow $h : j \rightarrow k$ such that $h \circ f = h \circ g$.

A κ -**directed preorder** is a preordered set that is κ -filtered when considered as a category; note that the third condition is then vacuous. A κ -**filtered diagram** (resp. κ -**directed diagram**) in a category \mathcal{C} is a functor $\mathbb{D} \rightarrow \mathcal{C}$ such that \mathbb{D} is a κ -filtered category (resp. κ -directed preorder). It is conventional to omit mention of κ when $\kappa = \aleph_0$.

Example 0.2.2. The category with one object $*$ and only one non-trivial arrow f is filtered if and only if $f = f \circ f$.

Example 0.2.3. Let X be any set. The set of all finite subsets of X , partially ordered by inclusion, is a directed preorder. More generally, if κ is any regular cardinal, then the set of all subsets of X with cardinality strictly less than κ is a κ -directed preorder.

Theorem 0.2.4. *Let \mathbf{U} be a pre-universe, let \mathbf{Set} be the category of \mathbf{U} -sets, and let κ be any regular cardinal. Given a \mathbf{U} -small category \mathbb{D} , the following are equivalent:*

- (i) \mathbb{D} is a κ -filtered category.
- (ii) The functor $\varinjlim_{\mathbb{D}} : [\mathbb{D}, \mathbf{Set}] \rightarrow \mathbf{Set}$ preserves limits for all diagrams that are both κ -small and \mathbf{U} -small.

Proof. The claim (i) \Rightarrow (ii) is very well known, and the converse is an exercise in using the Yoneda lemma and manipulating limits and colimits for diagrams of representable functors. \square

Definition 0.2.5. Let κ be a regular cardinal in a universe \mathbf{U}^+ and let \mathbf{U} be a pre-universe with $\mathbf{U} \subseteq \mathbf{U}^+$. A (κ, \mathbf{U}) -**compact object** in a locally \mathbf{U}^+ -small category \mathcal{C} is an object A such that the representable functor $\mathcal{C}(A, -) : \mathcal{C} \rightarrow \mathbf{Set}^+$ preserves colimits for all \mathbf{U} -small κ -filtered diagrams. A κ -**compact object** is one that is (κ, \mathbf{U}) -compact for all pre-universes \mathbf{U} .

Though the above definition is stated using a pre-universe \mathbf{U} contained in a universe \mathbf{U}^+ , the following lemma shows there is no dependence on \mathbf{U}^+ .

Lemma 0.2.6. *Let A be an object in a locally \mathbf{U}^+ -small category \mathcal{C} . The following are equivalent:*

- (i) A is a (κ, \mathbf{U}) -compact object in \mathcal{C} .

- (ii) For all \mathbf{U} -small κ -filtered diagrams $B : \mathbb{D} \rightarrow \mathcal{C}$, if $\lambda : B \Rightarrow \Delta C$ is a colimiting cocone, then for any morphism $f : A \rightarrow C$, there exist an object i in \mathbb{D} and a morphism $f' : A \rightarrow Bi$ in \mathcal{C} such that $f = \lambda_i \circ f'$; and moreover if $f = \lambda_j \circ f''$ for some morphism $f'' : A \rightarrow Bj$ in \mathcal{C} , then there exists an object k and a pair of arrows $g : i \rightarrow k, h : i \rightarrow k$ in \mathbb{D} such that $Bg \circ f' = Bh \circ f''$.

Proof. Use the explicit description of $\varinjlim_{\mathbb{D}} C(A, B)$ as a filtered colimit of sets; see Definition 1.1 in [LPAC], or Proposition 5.1.3 in [Borceux, 1994b]. \square

Corollary 0.2.7. Let $B : \mathbb{D} \rightarrow \mathcal{C}$ be a \mathbf{U} -small κ -filtered diagram, and let $\lambda : B \Rightarrow \Delta C$ be a colimiting cocone in \mathcal{C} . If C is a (κ, \mathbf{U}) -compact object in \mathcal{C} , then for some object i in \mathbb{D} , $\lambda_i : Bi \rightarrow C$ is a split epimorphism. \blacksquare

Lemma 0.2.8. Let A be an object in a category \mathcal{C} .

- (i) If \mathbf{U} is a pre-universe contained in a universe \mathbf{U}^+ and κ is a regular cardinal such that A is (κ, \mathbf{U}^+) -compact, then A is (κ, \mathbf{U}) -compact as well.
- (ii) If κ is a regular cardinal such that A is (κ, \mathbf{U}) -compact and λ is any regular cardinal such that $\kappa \leq \lambda$, then A is also (λ, \mathbf{U}) -compact.

Proof. Obvious. \blacksquare

Lemma 0.2.9. Let λ be a regular cardinal in a universe \mathbf{U}^+ , and let \mathbf{U} be a pre-universe with $\mathbf{U} \subseteq \mathbf{U}^+$. If $B : \mathbb{D} \rightarrow \mathcal{C}$ is a λ -small diagram of (λ, \mathbf{U}) -compact objects in a locally \mathbf{U}^+ -small category, then the colimit $\varinjlim_{\mathbb{D}} B$, if it exists, is a (λ, \mathbf{U}) -compact object in \mathcal{C} .

Proof. Use theorem 0.2.4 and the fact that $C(-, C) : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}^+$ maps colimits in \mathcal{C} to limits in \mathbf{Set}^+ . \blacksquare

Corollary 0.2.10. A retract of a (λ, \mathbf{U}) -compact object is also a (κ, \mathbf{U}) -compact object.

Proof. Suppose $r : A \rightarrow B$ and $s : B \rightarrow A$ are morphisms in \mathcal{C} such that $r \circ s = \text{id}_B$. Then $e = s \circ r$ is an idempotent morphism and the diagram below

$$A \begin{array}{c} \xrightarrow{\text{id}_A} \\ \xrightarrow[e]{} \end{array} A \xrightarrow{r} B$$

is a (split) coequaliser diagram in \mathcal{C} , so B is (λ, \mathbf{U}) -compact if A is. \blacksquare

Proposition 0.2.11. *Let \mathbf{U} be a pre-universe and let \mathbf{Set} be the category of \mathbf{U} -sets. For any \mathbf{U} -set A , the following are equivalent:*

- (i) *A has cardinality less than κ .*
- (ii) *The representable functor $\mathbf{Set}(A, -) : \mathbf{Set} \rightarrow \mathbf{Set}$ preserves colimits for all \mathbf{U} -small κ -filtered diagrams.*
- (iii) *The representable functor $\mathbf{Set}(A, -) : \mathbf{Set} \rightarrow \mathbf{Set}$ preserves colimits for all \mathbf{U} -small κ -directed diagrams.*

Proof. The claim (i) \Rightarrow (ii) follows from the theorem, and (ii) \Rightarrow (iii) is obvious. To see (iii) \Rightarrow (i), we may use [corollary 0.2.7](#) and the fact that every set is the directed union of its subsets of cardinality at most κ . ■

Corollary 0.2.12. *A set is κ -compact if and only if its cardinality is $< \kappa$.* ■

Definition 0.2.13. Let κ be a regular cardinal in a universe \mathbf{U} . A κ -**accessible \mathbf{U} -category** is a locally \mathbf{U} -small category \mathcal{C} satisfying the following conditions:

- \mathcal{C} has colimits for all \mathbf{U} -small κ -filtered diagrams.
- There exists a \mathbf{U} -set \mathcal{G} whose members are (κ, \mathbf{U}) -compact objects in \mathcal{C} such that, for every object B in \mathcal{C} , there exists a \mathbf{U} -small κ -filtered diagram of objects in \mathcal{G} with B as its colimit in \mathcal{C} .

We write $\mathbf{K}_\kappa^\mathbf{U}(\mathcal{C})$ for the full subcategory of \mathcal{C} spanned by the (κ, \mathbf{U}) -compact objects.

Example 0.2.14. The category of \mathbf{U} -sets is a κ -accessible \mathbf{U} -category for any regular cardinal κ in \mathbf{U} .

Remark 0.2.15. [Lemma 0.2.9](#) implies that, for each object A in an accessible \mathbf{U} -category, there exists a regular cardinal λ in \mathbf{U} such that A is (λ, \mathbf{U}) -compact.

Theorem 0.2.16. *Let \mathcal{C} be a locally \mathbf{U} -small category, and let κ be a regular cardinal in \mathbf{U} . There exist a locally \mathbf{U} -small category $\mathbf{Ind}_\mathbf{U}^\kappa(\mathcal{C})$ and a functor $\gamma : \mathcal{C} \rightarrow \mathbf{Ind}_\mathbf{U}^\kappa(\mathcal{C})$ with the following properties:*

- (i) *The objects of $\mathbf{Ind}_\mathbf{U}^\kappa(\mathcal{C})$ are \mathbf{U} -small κ -filtered diagrams $B : \mathbb{D} \rightarrow \mathcal{C}$, and γ sends an object C in \mathcal{C} to the corresponding trivial diagram $\mathbb{1} \rightarrow \mathcal{C}$ with value C .*

- (ii) *The functor $\gamma : \mathcal{C} \rightarrow \mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ is fully faithful, injective on objects, preserves all limits that exist in \mathcal{C} , and preserves all κ -small colimits that exist in \mathcal{C} .*
- (iii) *$\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ has colimits for all \mathbf{U} -small κ -filtered diagrams.*
- (iv) *For every object C in \mathcal{C} , the object γC is (κ, \mathbf{U}) -compact in $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$, and for each \mathbf{U} -small κ -filtered diagram $B : \mathbb{D} \rightarrow \mathcal{C}$, there is a canonical colimiting cocone $\gamma B \Rightarrow \Delta B$ in $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$.*
- (v) *If \mathcal{D} is a category with colimits for all \mathbf{U} -small κ -filtered diagrams, then for each functor $F : \mathcal{C} \rightarrow \mathcal{D}$, there exists a functor $\bar{F} : \mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C}) \rightarrow \mathcal{D}$ that preserves colimits for all \mathbf{U} -small κ -filtered diagrams in $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ such that $\gamma \bar{F} = F$, and given any functor $\bar{G} : \mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C}) \rightarrow \mathcal{D}$ whatsoever, the induced map $\text{Nat}(\bar{F}, \bar{G}) \rightarrow \text{Nat}(F, \gamma \bar{G})$ is a bijection.*

The category $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ is called the **free (κ, \mathbf{U}) -ind-completion** of \mathcal{C} , or the **category of (κ, \mathbf{U}) -ind-objects** in \mathcal{C} .

Proof. If $B : \mathbb{D} \rightarrow \mathcal{C}$ and $B' : \mathbb{D}' \rightarrow \mathcal{C}$ are two \mathbf{U} -small κ -filtered diagrams, then properties (ii) and (iii) together imply that

$$\text{Hom}(B', B) \cong \lim_{\leftarrow \mathbb{D}'} \lim_{\rightarrow \mathbb{D}} C(B', B)$$

and so, taking the RHS as the *definition* of the LHS, we need only find a suitable notion of composition to make $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ into a locally \mathbf{U} -small category. However, we observe that, if $N : \mathcal{C} \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is the Yoneda embedding, then

$$\text{Hom}\left(\lim_{\leftarrow \mathbb{D}'} NB', \lim_{\rightarrow \mathbb{D}} NB\right) \cong \lim_{\leftarrow \mathbb{D}'} \lim_{\rightarrow \mathbb{D}} C(B', B)$$

and, assuming property (v), the Yoneda embedding $N : \mathcal{C} \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ must extend along γ to a functor $\bar{N} : \mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C}) \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ that preserves colimits for \mathbf{U} -small κ -filtered diagram, so, in consideration of properties (i) and (iv), we may as well *define* the composition in $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ so that \bar{N} becomes fully faithful. This completes the definition of $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ as a category.

It remains to be shown that $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ actually has properties (ii), (iii), (iv), and (v); see Corollary 6.4.14 in [Borceux, 1994a] and Theorem 2.26 in [LPAC]. Note that the fact that γ preserves colimits for κ -small diagrams essentially follows from [theorem 0.2.4](#). \square

Proposition 0.2.17. *Let \mathbb{B} be a \mathbf{U} -small category and let κ be a regular cardinal in \mathbf{U} .*

- (i) $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B})$ is a κ -accessible \mathbf{U} -category.
- (ii) Every (κ, \mathbf{U}) -compact object in $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B})$ is a retract of an object of the form γB , where $\gamma : \mathbb{B} \rightarrow \mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B})$ is the canonical embedding.
- (iii) $\mathbf{K}_{\kappa}^{\mathbf{U}}(\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B}))$ is an essentially \mathbf{U} -small category.

Proof. (i). This claim more-or-less follows from the properties of $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B})$ explained in the previous theorem.

(ii). Use [corollary 0.2.10](#).

(iii). Since \mathbb{B} is \mathbf{U} -small and $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B})$ is locally \mathbf{U} -small, claim (ii) implies that $\mathbf{K}_{\kappa}^{\mathbf{U}}(\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B}))$ must be essentially \mathbf{U} -small. ■

Definition 0.2.18. Let κ be a regular cardinal in a universe \mathbf{U} . A (κ, \mathbf{U}) -**accessible functor** is a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ such that

- \mathcal{C} is a κ -accessible \mathbf{U} -category, and
- F preserves all colimits for \mathbf{U} -small κ -filtered diagrams.

We write $\mathbf{Acc}_{\kappa}^{\mathbf{U}}(\mathcal{C}, \mathcal{D})$ for the full subcategory of the functor category $[\mathcal{C}, \mathcal{D}]$ spanned by the (κ, \mathbf{U}) -accessible functors. An **accessible functor** is a functor that is (κ, \mathbf{U}) -accessible functor for some regular cardinal κ in some universe \mathbf{U} .

Theorem 0.2.19 (Classification of accessible categories). *Let κ be a regular cardinal in a universe \mathbf{U} , and let \mathcal{C} be a locally \mathbf{U} -small category. The following are equivalent:*

- (i) \mathcal{C} is a κ -accessible \mathbf{U} -category.
- (ii) The inclusion $\mathbf{K}_{\kappa}^{\mathbf{U}}(\mathcal{C}) \hookrightarrow \mathcal{C}$ extends along the embedding $\gamma : \mathcal{C} \rightarrow \mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathcal{C})$ to a (κ, \mathbf{U}) -accessible functor $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbf{K}_{\kappa}^{\mathbf{U}}(\mathcal{C})) \rightarrow \mathcal{C}$ that is fully faithful and essentially surjective on objects.
- (iii) There exist a \mathbf{U} -small category \mathbb{B} and a functor $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B}) \rightarrow \mathcal{C}$ that is fully faithful and essentially surjective on objects.

Proof. See Theorem 2.26 in [LPAC], or Theorem 5.35 in [Borceux, 1994b]. \square

Corollary 0.2.20. *If \mathcal{C} is a κ -accessible \mathbf{U} -category and \mathcal{D} is any category, then:*

- (i) *The restriction $\mathbf{Acc}_\kappa^{\mathbf{U}}(\mathcal{C}, \mathcal{D}) \rightarrow [\mathbf{K}_\kappa^{\mathbf{U}}(\mathcal{C}), \mathcal{D}]$ is fully faithful and surjective on objects.*
- (ii) *In particular, if \mathcal{D} is also locally \mathbf{U} -small, then $\mathbf{Acc}_\kappa^{\mathbf{U}}(\mathcal{C}, \mathcal{D})$ is equivalent to a locally \mathbf{U} -small category.*
- (iii) *If \mathcal{D} has colimits for all \mathbf{U} -small κ -filtered diagrams, then the inclusion $\mathbf{Acc}_\kappa^{\mathbf{U}}(\mathcal{C}, \mathcal{D}) \hookrightarrow [\mathcal{C}, \mathcal{D}]$ has a left adjoint. \blacksquare*

Definition 0.2.21. Let κ be a regular cardinal in a universe \mathbf{U} . A **locally κ -presentable \mathbf{U} -category** is a κ -accessible \mathbf{U} -category that is also \mathbf{U} -cocomplete. A **locally presentable \mathbf{U} -category** is one that is a locally κ -presentable \mathbf{U} -category for some regular cardinal κ in \mathbf{U} , and we often say ‘locally finitely presentable’ instead of ‘locally \aleph_0 -presentable’.

Example 0.2.22. The category of \mathbf{U} -sets is a locally κ -presentable \mathbf{U} -category for any regular cardinal κ in \mathbf{U} .

Lemma 0.2.23. *Let \mathcal{C} be a locally κ -presentable \mathbf{U} -category.*

- (i) *For any regular cardinal λ in \mathbf{U} , if $\kappa \leq \lambda$, then \mathcal{C} is a locally λ -presentable \mathbf{U} -category.*
- (ii) *With λ as above, if $F : \mathcal{C} \rightarrow \mathcal{D}$ is a (κ, \mathbf{U}) -accessible functor, then it is also a (λ, \mathbf{U}) -accessible functor.*
- (iii) *If \mathbf{U}^+ is any universe with $\mathbf{U} \in \mathbf{U}^+$, and \mathcal{C} is a locally κ -presentable \mathbf{U}^+ -category, then \mathcal{C} must be a preorder. \blacksquare*

Proof. (i). See the remark after Theorem 1.20 in [LPAC], or Propositions 5.3.2 and 5.2.3 in [Borceux, 1994b].

(ii). A λ -filtered diagram is certainly κ -filtered, so if F preserves colimits for all \mathbf{U} -small κ -filtered diagrams in \mathcal{C} , it must also preserve colimits for all \mathbf{U} -small λ -filtered diagrams.

(iii). This is a corollary of [theorem 0.1.10](#). \blacksquare

Corollary 0.2.24. *A category C is a locally presentable \mathbf{U} -category for at most one universe \mathbf{U} , provided C is not a preorder.*

Proof. Use [proposition 0.1.35](#) together with the above lemma. ■

Theorem 0.2.25 (Classification of locally presentable categories). *Let κ be a regular cardinal in a universe \mathbf{U} , let \mathbf{Set} be the category of \mathbf{U} -sets, and let C be a locally \mathbf{U} -small category. The following are equivalent:*

- (i) C is a locally κ -presentable \mathbf{U} -category.
- (ii) There exist a \mathbf{U} -small category \mathbb{B} that has colimits for κ -small diagrams and a functor $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B}) \rightarrow C$ that is fully faithful and essentially surjective on objects.
- (iii) The restricted Yoneda embedding $C \rightarrow [\mathbf{K}_{\kappa}^{\mathbf{U}}(C)^{\text{op}}, \mathbf{Set}]$ is fully faithful, (κ, \mathbf{U}) -accessible, and has a left adjoint.
- (iv) There exists a \mathbf{U} -small category \mathbb{A} and a fully faithful (κ, \mathbf{U}) -accessible functor $R : C \rightarrow [\mathbb{A}, \mathbf{Set}]$ such that \mathbb{A} has limits for all κ -small diagrams, R has a left adjoint, and R is essentially surjective onto the full subcategory of functors $\mathbb{A} \rightarrow \mathbf{Set}$ that preserve finite limits.
- (v) There exists a \mathbf{U} -small category \mathbb{A} and a fully faithful (κ, \mathbf{U}) -accessible functor $R : C \rightarrow [\mathbb{A}, \mathbf{Set}]$ such that R has a left adjoint.
- (vi) C is a κ -accessible \mathbf{U} -category and is \mathbf{U} -complete.

Proof. See [Proposition 1.27](#), [Corollary 1.28](#), [Theorem 1.46](#), and [Corollary 2.47](#) in [\[LPAC\]](#), or [Theorems 5.2.7](#) and [5.5.8](#) in [\[Borceux, 1994b\]](#). □

Remark 0.2.26. If C is equivalent to $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B})$ for some \mathbf{U} -small category \mathbb{B} that has limits for all κ -small diagrams, then \mathbb{B} must be equivalent to $\mathbf{K}_{\kappa}^{\mathbf{U}}(C)$ by [proposition 0.2.17](#). In other words, every locally κ -presentable \mathbf{U} -category is, up to equivalence, the (κ, \mathbf{U}) -ind-completion of an essentially unique \mathbf{U} -small κ -complete category.

Example 0.2.27. Obviously, for any \mathbf{U} -small category \mathbb{A} , the functor category $[\mathbb{A}, \mathbf{Set}]$ is locally finitely presentable. More generally, one may show that for any κ -ary algebraic theory \mathbb{T} , possibly many-sorted, the category of \mathbb{T} -algebras in \mathbf{U} is a locally κ -presentable \mathbf{U} -category. The above theorem can also be used to

show that \mathbf{Cat} , the category of \mathbf{U} -small categories, is a locally finitely presentable \mathbf{U} -small category.

Corollary 0.2.28. *Let C be a locally κ -presentable \mathbf{U} -category. For any \mathbf{U} -small κ -filtered diagram \mathbb{D} , $\varinjlim_{\mathbb{D}} : [\mathbb{D}, C] \rightarrow C$ preserves κ -small limits.*

Proof. The claim is certainly true when $C = [\mathbb{A}, \mathbf{Set}]$, by [theorem 0.2.4](#). In general, choose a (κ, \mathbf{U}) -accessible fully faithful functor $R : C \rightarrow [\mathbb{A}, \mathbf{Set}]$ with a left adjoint, and simply note that R creates limits for all \mathbf{U} -small diagrams as well as colimits for all \mathbf{U} -small κ -filtered diagrams. \blacksquare

Proposition 0.2.29. *If C is a locally κ -presentable \mathbf{U} -category and \mathbb{D} is any \mathbf{U} -small category, then the functor category $[\mathbb{D}, C]$ is also a locally κ -presentable category.*

Proof. This can be proven using the classification theorem by noting that the 2-functor $[\mathbb{D}, -]$ preserves reflective subcategories, but see also [Corollary 1.54](#) in [\[LPAC\]](#). \square

It is commonplace to say ‘ λ -presentable object’ instead of ‘ λ -compact object’, especially in algebraic contexts. The following proposition justifies the alternative terminology:

Proposition 0.2.30. *Let C be a locally κ -presentable \mathbf{U} -category, and let λ be a regular cardinal in \mathbf{U} with $\lambda \geq \kappa$. If \mathcal{H} is a small full subcategory of C such that*

- *every (κ, \mathbf{U}) -compact object in C is isomorphic to an object in \mathcal{H} , and*
- *\mathcal{H} is closed in C under colimits for λ -small diagrams,*

then every (λ, \mathbf{U}) -compact object in C is isomorphic to an object in \mathcal{H} . In particular, $\mathbf{K}_\kappa^{\mathbf{U}}(C)$ is the smallest replete full subcategory of C containing $\mathbf{K}_\kappa^{\mathbf{U}}(C)$ and closed in C under colimits for λ -small diagrams.

Proof. Let C be any (λ, \mathbf{U}) -compact object in C . Clearly, the comma category $(\mathcal{H} \downarrow C)$ is a \mathbf{U} -small λ -filtered category. Let $\mathcal{G} = \mathcal{H} \cap \mathbf{K}_\kappa^{\mathbf{U}}(C)$. One can show that $(\mathcal{G} \downarrow C)$ is a cofinal subcategory in $(\mathcal{H} \downarrow C)$, and the classification theorem ([0.2.25](#)) implies that the tautological cocone on the diagram $(\mathcal{G} \downarrow C) \rightarrow C$ is colimiting, so the tautological cocone on the diagram $(\mathcal{H} \downarrow C) \rightarrow C$ is also colimiting. Now, by [corollary 0.2.7](#), C is a retract of an object in \mathcal{H} , and hence C must be isomorphic to an object in \mathcal{H} , because \mathcal{H} is closed under coequalisers.

TODO: Simplify this argument.

For the final claim, note that $\mathbf{K}_\lambda^{\mathbf{U}}(C)$ is certainly a replete full subcategory of C and contained in any replete full subcategory containing $\mathbf{K}_\kappa^{\mathbf{U}}(C)$ and closed in C under colimits for λ -small diagrams, so we just have to show that $\mathbf{K}_\lambda^{\mathbf{U}}(C)$ is also closed in C under colimits for λ -small diagrams; for this, we simply appeal to [lemma 0.2.9](#). \blacksquare

Proposition 0.2.31. *Let C be a locally κ -presentable \mathbf{U} -category and let \mathbb{D} be a μ -small category in \mathbf{U} . The (λ, \mathbf{U}) -compact objects in $[\mathbb{D}, C]$ are precisely the diagrams $\mathbb{D} \rightarrow C$ that are componentwise (λ, \mathbf{U}) -compact, so long as $\lambda \geq \max\{\kappa, \mu\}$.*

Proof. First, note that Mac Lane's subdivision category^[1] \mathbb{D}^\S is also μ -small, so $[\mathbb{D}, C](A, B)$ is computed as the limit of a μ -small diagram of hom-sets. More precisely, using end notation,

$$[\mathbb{D}, C](A, B) = \int_{d:\mathbb{D}} C(Ad, Bd)$$

and so if A is componentwise (λ, \mathbf{U}) -compact, then $[\mathbb{D}, C](A, -)$ preserves colimits for \mathbf{U} -small λ -filtered diagrams, hence A is itself (λ, \mathbf{U}) -compact.

Now, suppose A is a (λ, \mathbf{U}) -compact object in $[\mathbb{D}, C]$. Let d be an object in \mathbb{D} , let $d^* : [\mathbb{D}, C] \rightarrow C$ be evaluation at d , and let $d_* : C \rightarrow [\mathbb{D}, C]$ be the right adjoint, which is explicitly given by

$$(d_*C)(d') = \mathbb{D}(d', d) \pitchfork C$$

where \pitchfork is defined by following adjunction:

$$\mathbf{Set}(X, C(C, C')) \cong C(C, X \pitchfork C')$$

The unit $\eta_A : A \rightarrow d_*d^*A$ is constructed using the universal property of \pitchfork in the obvious way, and the counit $\varepsilon_C : d^*d_*C \rightarrow C$ is the projection $\mathbb{D}(d, d) \pitchfork C \rightarrow C$ corresponding to $\text{id}_d \in \mathbb{D}(d, d)$. Since C is a locally λ -presentable \mathbf{U} -category, there exist a \mathbf{U} -small λ -filtered diagram $B : \mathbb{J} \rightarrow C$ consisting of (λ, \mathbf{U}) -compact objects in C and a colimiting cocone $\alpha : B \Rightarrow \Delta d^*A$, and since each $\mathbb{D}(d', d)$ has cardinality less than μ , the cocone $d_*\alpha : d_*B \Rightarrow \Delta d_*d^*A$ is also colimiting, by [corollary 0.2.28](#). [Lemma 0.2.6](#) then implies $\eta_A : A \rightarrow d_*d^*A$ factors through $d_*\alpha_j : d_*(Bj) \rightarrow d_*d^*A$ for some j in \mathbb{J} , say

$$\eta_A = d_*\alpha_j \circ \sigma$$

^[1] See [CWM, Ch. IX, § 5].

for some $\sigma : A \rightarrow d_* B_j$. But then, by the triangle identity,

$$\text{id}_{Ad} = \varepsilon_{Ad} \circ d^* \eta_A = \varepsilon_{Ad} \circ d^* d_* \alpha_j \circ d^* \sigma = \alpha_j \circ \varepsilon_{B_j} \circ d^* \sigma$$

and so $\alpha_j : B_j \rightarrow Ad$ is a split epimorphism, hence Ad is a (λ, \mathbf{U}) -compact object, by [corollary 0.2.10](#). ■

Remark 0.2.32. The claim in the above proposition can fail if $\mu > \lambda \geq \kappa$. For example, we could take $\mathcal{C} = \mathbf{Set}$, with \mathbb{D} being the ordinal ω considered as a category; then the terminal object in $[\mathbb{D}, \mathbf{Set}]$ is componentwise finite, but is not itself an \aleph_0 -compact object in \mathbf{Set} .

Lemma 0.2.33. *Let κ and λ be regular cardinals in a universe \mathbf{U} , with $\kappa \leq \lambda$.*

- (i) *If \mathcal{D} is a locally λ -presentable \mathbf{U} -category, \mathcal{C} is a locally \mathbf{U} -small category, and $G : \mathcal{D} \rightarrow \mathcal{C}$ is a (λ, \mathbf{U}) -accessible functor that preserves limits for all \mathbf{U} -small diagrams in \mathcal{C} , then, for any (κ, \mathbf{U}) -compact object C in \mathcal{C} , the comma category $(C \downarrow G)$ has an initial object.*
- (ii) *If \mathcal{C} is a locally κ -presentable \mathbf{U} -category, \mathcal{D} is a locally \mathbf{U} -small category, and $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor that preserves colimits for all \mathbf{U} -small diagrams in \mathcal{C} , then, for any object D in \mathcal{D} , the comma category $(F \downarrow D)$ has a terminal object.*

Proof. (i). Let \mathcal{F} be the full subcategory of $(C \downarrow G)$ spanned by those (D, g) where D is a (λ, \mathbf{U}) -compact object in \mathcal{D} . G preserves colimits for all \mathbf{U} -small λ -filtered diagrams, so, by [lemma 0.2.6](#), \mathcal{F} must be a weakly initial family in $(C \downarrow G)$. [Proposition 0.2.17](#) implies \mathcal{F} is an essentially \mathbf{U} -small category, and since \mathcal{D} has limits for all \mathbf{U} -small diagrams and G preserves them, $(C \downarrow G)$ is also \mathbf{U} -complete. Thus, the inclusion $\mathcal{F} \hookrightarrow (C \downarrow G)$ has a limit, and it can be shown that this is an initial object in $(C \downarrow G)$.^[2]

(ii). Let \mathcal{G} be the full subcategory of $(F \downarrow D)$ spanned by those (C, f) where C is a (κ, \mathbf{U}) -compact object in \mathcal{C} ; note that [proposition 0.2.17](#) implies \mathcal{G} is an essentially \mathbf{U} -small category. Since \mathcal{C} has colimits for all \mathbf{U} -small diagrams and F preserves them, $(F \downarrow D)$ is also \mathbf{U} -cocomplete.^[3] Let (C, f) be a colimit for the inclusion $\mathcal{G} \hookrightarrow (F \downarrow D)$. It is not hard to check that (C, f) is a weakly terminal

^[2] See Theorem 1 in [\[CWM, Ch. X, § 2\]](#).

^[3] See the Lemma in [\[CWM, Ch. V, § 6\]](#).

object in $(F \downarrow D)$, so the formal dual of Freyd's initial object lemma^[4] gives us a terminal object in $(F \downarrow D)$; explicitly, it may be constructed as the joint coequaliser of all the endomorphisms of (C, f) . ■

Theorem 0.2.34 (Accessible adjoint functor theorem). *Let κ and λ be regular cardinals in a universe \mathbf{U} , with $\kappa \leq \lambda$, let \mathcal{C} be a locally κ -presentable \mathbf{U} -category, and let \mathcal{D} be a locally λ -presentable \mathbf{U} -category.*

Given a functor $F : \mathcal{C} \rightarrow \mathcal{D}$, the following are equivalent:

- (i) *F has a right adjoint $G : \mathcal{D} \rightarrow \mathcal{C}$, and G is a (λ, \mathbf{U}) -accessible functor.*
- (ii) *F preserves colimits for all \mathbf{U} -small diagrams and sends (κ, \mathbf{U}) -compact objects in \mathcal{C} to (λ, \mathbf{U}) -compact objects in \mathcal{D} .*
- (iii) *F has a right adjoint and sends (κ, \mathbf{U}) -compact objects in \mathcal{C} to (λ, \mathbf{U}) -compact objects in \mathcal{D} .*

On the other hand, given a functor $G : \mathcal{D} \rightarrow \mathcal{C}$, the following are equivalent:

- (iv) *G has a left adjoint $F : \mathcal{C} \rightarrow \mathcal{D}$, and F sends (κ, \mathbf{U}) -compact objects in \mathcal{C} to (λ, \mathbf{U}) -compact objects in \mathcal{D} .*
- (v) *G is a (λ, \mathbf{U}) -accessible functor and preserves limits for all \mathbf{U} -small diagrams.*
- (vi) *G is a (λ, \mathbf{U}) -accessible functor and there exist a functor $F_0 : \mathbf{K}_\kappa^{\mathbf{U}}(\mathcal{C}) \rightarrow \mathcal{D}$ and hom-set bijections*

$$C(\mathcal{C}, GD) \cong D(F_0\mathcal{C}, D)$$

natural in D for each (κ, \mathbf{U}) -compact object C in \mathcal{C} , where D varies in \mathcal{D} .

Proof. We will need to refer back to the details of the proof of this theorem later, so here is a sketch of the constructions involved.

(i) \Rightarrow (ii). If F is a left adjoint, then F certainly preserves colimits for all \mathbf{U} -small diagrams. Given a (κ, \mathbf{U}) -compact object C in \mathcal{C} and a \mathbf{U} -small λ -filtered diagram

^[4] See Theorem I in [CWM, Ch. V, § 6].

$B : \mathbb{J} \rightarrow \mathcal{D}$, observe that

$$\begin{aligned} \mathcal{D}\left(FC, \varinjlim_{\mathbb{J}} B\right) &\cong \mathcal{C}\left(C, G \varinjlim_{\mathbb{J}} B\right) \cong \mathcal{C}\left(C, \varinjlim_{\mathbb{J}} GB\right) \\ &\cong \varinjlim_{\mathbb{J}} \mathcal{C}(C, GB) \cong \varinjlim_{\mathbb{J}} \mathcal{C}(FC, B) \end{aligned}$$

and thus FC is indeed a (λ, \mathbf{U}) -compact object in \mathcal{D} .

(ii) \Rightarrow (iii). It is enough to show that, for each object D in \mathcal{D} , the comma category $(F \downarrow D)$ has a terminal object (GD, ε_D) ; ^[5] but this was done in the previous lemma.

(iii) \Rightarrow (i). Given a (κ, \mathbf{U}) -compact object C in \mathcal{C} and a \mathbf{U} -small λ -filtered diagram $B : \mathbb{J} \rightarrow \mathcal{D}$, observe that

$$\begin{aligned} \mathcal{C}\left(C, G \varinjlim_{\mathbb{J}} B\right) &\cong \mathcal{D}\left(FC, \varinjlim_{\mathbb{J}} B\right) \cong \varinjlim_{\mathbb{J}} \mathcal{C}(FC, B) \\ &\cong \varinjlim_{\mathbb{J}} \mathcal{C}(C, GB) \cong \mathcal{C}\left(C, \varinjlim_{\mathbb{J}} GB\right) \end{aligned}$$

because FC is a (λ, \mathbf{U}) -compact object in \mathcal{D} ; but [theorem 0.2.25](#) says the restricted Yoneda embedding $\mathcal{C} \rightarrow [\mathbf{K}_{\kappa}^{\mathbf{U}}(\mathcal{C})^{\text{op}}, \mathbf{Set}]$ is fully faithful, so this is enough to conclude that G preserves colimits for \mathbf{U} -small λ -filtered diagrams.

(iv) \Rightarrow (v). If G is a right adjoint, then G certainly preserves colimits for all \mathbf{U} -small diagrams; the rest of the claim is subsumed by (iii) \Rightarrow (i).

(v) \Rightarrow (vi). It is enough to show that, for each (κ, \mathbf{U}) -compact object C in \mathcal{C} , the comma category $(C \downarrow G)$ has an initial object (F_0C, η_C) ; but this was done in the previous lemma. It is clear how to make F_0 into a functor $\mathbf{K}_{\kappa}^{\mathbf{U}}(\mathcal{C}) \rightarrow \mathcal{D}$.

(vi) \Rightarrow (iv). We use [theorems 0.2.16](#) and [0.2.25](#) to extend $F_0 : \mathbf{K}_{\kappa}^{\mathbf{U}}(\mathcal{C}) \rightarrow \mathcal{D}$ along the inclusion $\mathbf{K}_{\kappa}^{\mathbf{U}}(\mathcal{C}) \hookrightarrow \mathcal{C}$ to get (κ, \mathbf{U}) -accessible functor $F : \mathcal{C} \rightarrow \mathcal{D}$. We then observe that, for any \mathbf{U} -small κ -filtered diagram $A : \mathbb{I} \rightarrow \mathcal{C}$ of (κ, \mathbf{U}) -compact

^[5] See Theorem 2 in [\[CWM, Ch. IV, § 1\]](#).

objects in \mathcal{C} ,

$$\begin{aligned} \mathcal{C}\left(\varinjlim A, GD\right) &\cong \varprojlim \mathcal{C}(A, GD) \cong \varprojlim \mathcal{C}(F_0 A, D) \\ &\cong \mathcal{C}\left(\varinjlim F A, D\right) \cong \mathcal{C}\left(F \varinjlim A, D\right) \end{aligned}$$

is a series of bijections natural in D , where D varies in \mathcal{D} ; but \mathcal{C} is a locally κ -presentable \mathbf{U} -category, so this is enough to show that F is a left adjoint of G . The remainder of the claim is a corollary of (i) \Rightarrow (ii). \blacksquare

Corollary 0.2.35. *Let \mathcal{C} and \mathcal{D} be locally presentable \mathbf{U} -categories. If a functor $G : \mathcal{D} \rightarrow \mathcal{C}$ has a left adjoint, then there exists a regular cardinal μ in \mathbf{U} such that G is a (μ, \mathbf{U}) -accessible functor.*

Proof. Suppose \mathcal{C} is a locally κ -presentable \mathbf{U} -category, \mathcal{D} is a locally λ -presentable \mathbf{U} -category, and $F : \mathcal{C} \rightarrow \mathcal{D}$ is a left adjoint for G . Since $\mathbf{K}_\kappa^{\mathbf{U}}(\mathcal{C})$ is an essentially \mathbf{U} -small category, recalling [lemma 0.2.8](#), there certainly exists a regular cardinal μ in \mathbf{U} such that $\mu \geq \lambda$ and F sends (κ, \mathbf{U}) -compact objects in \mathcal{C} to (μ, \mathbf{U}) -compact objects in \mathcal{D} . The above theorem, plus [lemma 0.2.23](#), implies G is an (μ, \mathbf{U}) -accessible functor. \blacksquare

0.3 Change of universe

Prerequisites. §§ [0.1](#), [0.2](#).

Having introduced universes into our ontology, it becomes necessary to ask whether an object with some universal property retains that property when we enlarge the universe. Though it sounds inconceivable, there do exist examples of badly-behaved constructions that are not stable under change-of-universe; for example, Waterhouse [[1975](#)] defined a functor $F : \mathbf{CRing} \rightarrow \mathbf{Set}^+$, where \mathbf{CRing} is the category of commutative rings in a universe \mathbf{U} and \mathbf{Set}^+ is the category of \mathbf{U}^+ -sets for some universe \mathbf{U}^+ with $\mathbf{U} \in \mathbf{U}^+$, such that the value of F at any given commutative ring in \mathbf{U} does not depend on \mathbf{U} , and yet the value of the fpqc sheaf associated with F at the field \mathbb{Q} depends on the size of \mathbf{U} .

Many of the universal properties of interest concern adjunctions, so that is where we begin.

Definition 0.3.1. Let $F \dashv G : \mathcal{D} \rightarrow \mathcal{C}$ and $F' \dashv G' : \mathcal{D}' \rightarrow \mathcal{C}'$ be adjunctions, and let $H : \mathcal{C} \rightarrow \mathcal{C}'$ and $K : \mathcal{D} \rightarrow \mathcal{D}'$ be functors. The **mate** of a natural transformation $\alpha : HG \Rightarrow G'K$ is the natural transformation

$$\varepsilon' K F \bullet F' \alpha F \bullet F' H \eta : F' H \Rightarrow K F$$

where $\eta : \text{id}_{\mathcal{C}} \Rightarrow GF$ is the unit of $F \dashv G$ and $\varepsilon : F'G' \Rightarrow \text{id}_{\mathcal{D}'}$ is the counit of $F' \dashv G'$; dually, the **mate** of a natural transformation $\beta : F'H \Rightarrow KF$ is the natural transformation

$$G' K \varepsilon \bullet G' \beta G \bullet \eta' H G : H G \Rightarrow G' K$$

where $\eta' : \text{id}_{\mathcal{C}'} \Rightarrow G'F'$ is the unit of $F' \dashv G'$ and $\varepsilon : FG \Rightarrow \text{id}_{\mathcal{D}}$ is the counit of $F \dashv G$.

Lemma 0.3.2. *In the above notation, the two mates constructions constitute a mutually inverse pair of bijections*

$$\text{Nat}(F' H, K F) \cong \text{Nat}(H G, G' K)$$

and moreover, given a further adjunction $F'' \dashv G'' : \mathcal{C}'' \rightarrow \mathcal{D}''$ and functors $H' : \mathcal{C}' \rightarrow \mathcal{C}''$ and $K' : \mathcal{D}' \rightarrow \mathcal{D}''$, if $\alpha : HG \Rightarrow G'K$ and $\alpha' : H'G' \Rightarrow G''K'$ have mates $\beta : F'H \Rightarrow KF$ and $\beta' : F''H' \Rightarrow K'F'$ respectively, then the composite natural transformation $\alpha' K \bullet H' \alpha : H' H G \Rightarrow H'' K' K$ has mate $K' \beta \bullet \beta' H : F'' H' H \Rightarrow K' K F$.

Proof. This is an exercise in using the triangle identities for adjunctions. \square

Definition 0.3.3. Given a diagram of the form

$$\begin{array}{ccc} \mathcal{D} & \xrightarrow{K} & \mathcal{D}' \\ G \downarrow & \nearrow \alpha & \downarrow G' \\ \mathcal{C} & \xrightarrow{H} & \mathcal{C}' \end{array}$$

where $\alpha : HG \Rightarrow G'K$ is a natural isomorphism, $F \dashv G$ and $F' \dashv G'$, we say the diagram satisfies the **left Beck–Chevalley condition** if the mate of α is also a natural isomorphism. Dually, given a diagram of the form

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{H} & \mathcal{C}' \\ F \downarrow & \nwarrow \beta & \downarrow F' \\ \mathcal{D} & \xrightarrow{K} & \mathcal{D}' \end{array}$$

where $\beta : F'H \Rightarrow KF$ is a natural isomorphism, $F \dashv G$ and $F' \dashv G'$, we say the diagram satisfies the **right Beck–Chevalley condition** if the mate of β is also a natural isomorphism.

Remark 0.3.4. Unfortunately, the Beck–Chevalley conditions are not vacuous. For example, consider the following (strictly!) commutative diagram of forgetful functors:

$$\begin{array}{ccc} \mathbf{CRing} & \longrightarrow & \mathbf{Ab} \\ \downarrow & & \downarrow \\ \mathbf{Set} & \xrightarrow{\text{id}} & \mathbf{Set} \end{array}$$

The mate of the trivial natural transformation in the above diagram is the group homomorphism $\mathbb{Z}X \rightarrow \mathbb{Z}[X]$ that sends a generator in $\mathbb{Z}X$ to the corresponding generator in $\mathbb{Z}[X]$; clearly, this is never an isomorphism. However, this is unsurprising: we do not expect the additive group of free commutative ring generated by X to be naturally isomorphic to the free abelian group generated by X .

Example 0.3.5. Let \mathcal{C} be a category with pullbacks, and suppose

$$\begin{array}{ccc} Z & \xrightarrow{z} & X \\ g \downarrow & & \downarrow f \\ W & \xrightarrow{w} & Y \end{array}$$

is a pullback square in \mathcal{C} . Let $\Sigma_f : \mathcal{C}_{/X} \rightarrow \mathcal{C}_{/Y}$ etc. be the functor that sends an object $p : E \rightarrow X$ in $\mathcal{C}_{/X}$ to the object $f \circ p : E \rightarrow Y$ in $\mathcal{C}_{/Y}$, and consider the induced (strictly!) commutative diagram of functors:

$$\begin{array}{ccc} \mathcal{C}_{/Z} & \xrightarrow{\Sigma_z} & \mathcal{C}_{/X} \\ \Sigma_g \downarrow & & \downarrow \Sigma_f \\ \mathcal{C}_{/W} & \xrightarrow{\Sigma_w} & \mathcal{C}_{/Y} \end{array}$$

Since \mathcal{C} has pullbacks, Σ_g and Σ_f have right adjoints, and the pullback pasting lemma then implies that the above square satisfies the right Beck–Chevalley condition.

Lemma 0.3.6. *Given a diagram of the form*

$$\begin{array}{ccc} \mathcal{D} & \xrightarrow{K} & \mathcal{D}' \\ G \downarrow & \nearrow \alpha & \downarrow G' \\ \mathcal{C} & \xrightarrow{H} & \mathcal{C}' \end{array}$$

where $\alpha : HG \Rightarrow G'K$ is a natural isomorphism, $F \dashv G$ and $F' \dashv G'$, the diagram satisfies the left Beck–Chevalley condition if and only if, for every object C in \mathcal{C} , the functor $(C \downarrow G) \rightarrow (HC \downarrow G')$ sending an object (D, f) in the comma category $(C \downarrow G)$ to the object $(KD, \alpha_D \circ Hf)$ in $(HC \downarrow G')$ preserves initial objects.

Proof. We know (FC, η_C) is an initial object of $(C \downarrow G)$ and $(F'HC, \eta'_{HC})$ is an initial object of $(HC \downarrow G')$, so there is a unique morphism $\beta_C : F'HC \rightarrow KFC$ such that $G'\beta_C \circ \eta'_{HC} = \alpha_{FC} \circ H\eta_C$. However, we observe that

$$\begin{aligned} \beta_C &= \beta_C \circ \varepsilon'_{F'HC} \circ F'\eta'_{HC} \\ &= \varepsilon'_{KFC} \circ F'G'\beta_C \circ F'\eta'_{HC} \\ &= \varepsilon'_{KFC} \circ F'\alpha_{FC} \circ F'H\eta_C \end{aligned}$$

so β_C is precisely the component at C of the mate of α . Thus β_C is an isomorphism for all C if and only if the Beck–Chevalley condition holds. \blacksquare

Definition 0.3.7. Let κ be a regular cardinal in a universe \mathbf{U} , and let \mathbf{U}^+ be a universe with $\mathbf{U} \subseteq \mathbf{U}^+$. A $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -**accessible extension** is a (κ, \mathbf{U}) -accessible functor $i : \mathcal{C} \rightarrow \mathcal{C}^+$ such that

- \mathcal{C} is a κ -accessible \mathbf{U} -category,
- \mathcal{C}^+ is a κ -accessible \mathbf{U}^+ -category,
- i sends (κ, \mathbf{U}) -compact objects in \mathcal{C} to (κ, \mathbf{U}^+) -compact objects in \mathcal{C}^+ , and
- the functor $\mathbf{K}_\kappa^{\mathbf{U}}(\mathcal{C}) \rightarrow \mathbf{K}_\kappa^{\mathbf{U}^+}(\mathcal{C}^+)$ so induced by i is fully faithful and essentially surjective on objects.

Remark 0.3.8. Let \mathbb{B} be a \mathbf{U} -small category in which idempotents split. Then the (κ, \mathbf{U}) -accessible functor $\mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B}) \rightarrow \mathbf{Ind}_{\mathbf{U}^+}^{\kappa}(\mathbb{B})$ obtained by extending the embedding $\gamma^+ : \mathbb{B} \rightarrow \mathbf{Ind}_{\mathbf{U}^+}^{\kappa}(\mathbb{B})$ along $\gamma : \mathbb{B} \rightarrow \mathbf{Ind}_{\mathbf{U}}^{\kappa}(\mathbb{B})$ is a $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -extension, by [proposition 0.2.17](#). The classification theorem ([0.2.19](#)) implies all examples of $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -accessible extensions are essentially of this form.

Proposition 0.3.9. *Let $i : C \rightarrow C^+$ be a $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -accessible extension.*

- (i) *C is a locally κ -presentable \mathbf{U} -category if and only if C^+ is a locally κ -presentable \mathbf{U}^+ -category.*
- (ii) *The functor $i : C \rightarrow C^+$ is fully faithful.*
- (iii) *If $B : \mathbb{J} \rightarrow C$ is any diagram (not necessarily \mathbf{U} -small) and C has a limit for B , then i preserves this limit.*

Proof. (i). If C is a locally κ -presentable \mathbf{U} -category, then $\mathbf{K}_\kappa^{\mathbf{U}}(C)$ has colimits for all κ -small diagrams, so $\mathbf{K}_\kappa^{\mathbf{U}^+}(C^+)$ also has colimits for all κ -small diagrams. The classification theorem (0.2.19) then implies C^+ is a locally κ -presentable \mathbf{U}^+ -category. Reversing this argument proves the converse.

(ii). Let $A : \mathbb{I} \rightarrow C$ and $B : \mathbb{J} \rightarrow C$ be two \mathbf{U} -small κ -filtered diagrams of (κ, \mathbf{U}) -compact objects in C . Then,

$$\begin{aligned} C\left(\lim_{\mathbb{I}} A, \lim_{\mathbb{J}} B\right) &\cong \lim_{\mathbb{I}} \lim_{\mathbb{J}} C(A, B) \cong \lim_{\mathbb{I}} \lim_{\mathbb{J}} C^+(iA, iB) \\ &\cong C^+\left(\lim_{\mathbb{I}} iA, \lim_{\mathbb{J}} iB\right) \cong C^+\left(i \lim_{\mathbb{I}} A, i \lim_{\mathbb{J}} B\right) \end{aligned}$$

because i is (κ, \mathbf{U}) -accessible and is fully faithful on the subcategory $\mathbf{K}_\kappa^{\mathbf{U}}(C)$, and therefore $i : C \rightarrow C^+$ itself is fully faithful. Note that this hinges crucially on [theorem 0.1.30](#).

(iii). Let $B : \mathbb{J} \rightarrow C$ be any diagram. We observe that, for any (κ, \mathbf{U}) -compact object C in C ,

$$\begin{aligned} C^+\left(iC, i \lim_{\mathbb{J}} B\right) &\cong C\left(C, \lim_{\mathbb{J}} B\right) && \text{because } i \text{ is fully faithful} \\ &\cong \lim_{\mathbb{J}} C(C, B) && \text{by definition of limit} \\ &\cong \lim_{\mathbb{J}} C^+(iC, iB) && \text{because } i \text{ is fully faithful} \end{aligned}$$

but we know the restricted Yoneda embedding $C^+ \rightarrow [\mathbf{K}_\kappa^{\mathbf{U}}(C)^{\text{op}}, \mathbf{Set}^+]$ is fully faithful, so this is enough to conclude that $i \lim_{\mathbb{J}} B$ is the limit of iB in C^+ . ■

Remark 0.3.10. Similar methods show that any fully faithful functor $\mathcal{C} \rightarrow \mathcal{C}^+$ satisfying the four bulleted conditions in the definition above is necessarily (κ, \mathbf{U}) -accessible.

Lemma 0.3.11. *Let \mathbf{U} and \mathbf{U}^+ be universes, with $\mathbf{U} \in \mathbf{U}^+$, and let κ be a regular cardinal in \mathbf{U} . Suppose:*

- \mathcal{C} and \mathcal{D} are locally κ -presentable \mathbf{U} -categories.
- \mathcal{C}^+ and \mathcal{D}^+ are locally κ -presentable \mathbf{U}^+ -categories.
- $i : \mathcal{C} \rightarrow \mathcal{C}^+$ and $j : \mathcal{D} \rightarrow \mathcal{D}^+$ are $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -accessible extensions.

Given a strictly commutative diagram of the form below,

$$\begin{array}{ccc} \mathcal{D} & \xrightarrow{j} & \mathcal{D}^+ \\ G \downarrow & & \downarrow G^+ \\ \mathcal{C} & \xrightarrow{i} & \mathcal{C}^+ \end{array}$$

where G is (κ, \mathbf{U}) -accessible, G^+ is (κ, \mathbf{U}^+) -accessible, if both have left adjoints, then the diagram satisfies the left Beck–Chevalley condition.

Proof. Let C be a (κ, \mathbf{U}) -compact object in \mathcal{C} . Inspecting the proof of [theorem 0.2.34](#), we see that the functor $(\mathcal{C} \downarrow G) \rightarrow (i\mathcal{C} \downarrow G^+)$ induced by j preserves initial objects. As in the proof of [lemma 0.3.6](#), this implies the component at C of the left Beck–Chevalley natural transformation $F^+i \Rightarrow jF$ is an isomorphism; but C is generated by $\mathbf{K}_\kappa^{\mathbf{U}}(\mathcal{C})$ and the functors F, F^+, i, j all preserve colimits for \mathbf{U} -small κ -filtered diagrams, so in fact $F^+i \Rightarrow jF$ is a natural isomorphism. ■

Proposition 0.3.12. *If $i : \mathcal{C} \rightarrow \mathcal{C}^+$ is a $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -accessible extension and \mathcal{C} is a locally κ -presentable \mathbf{U} -category, then i preserves colimits for all \mathbf{U} -small diagrams in \mathcal{C} .*

Proof. It is well-known that a functor preserves colimits for all \mathbf{U} -small diagrams if and only if it preserves coequalisers for all parallel pairs and coproducts for all \mathbf{U} -small families, but coproducts for \mathbf{U} -small families can be constructed in a uniform way using coproducts for κ -small families and colimits for \mathbf{U} -small κ -filtered diagrams. It is therefore enough to show that $i : \mathcal{C} \rightarrow \mathcal{C}^+$ preserves all colimits for κ -small diagrams, since i is already (κ, \mathbf{U}) -accessible.

Let \mathbb{D} be a κ -small category. Recalling [proposition 0.1.12](#), our problem amounts to showing that the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{i} & \mathcal{C}^+ \\ \Delta \downarrow & & \downarrow \Delta^+ \\ [\mathbb{D}, \mathcal{C}] & \xrightarrow{i_*} & [\mathbb{D}, \mathcal{C}^+] \end{array}$$

satisfies the left Beck–Chevalley condition. It is clear that i_* is fully faithful. Colimits for \mathbf{U} -small diagrams in $[\mathbb{D}, \mathcal{C}]$ and in $[\mathbb{D}, \mathcal{C}^+]$ are computed componentwise, so Δ and i_* are certainly (κ, \mathbf{U}) -accessible, and Δ^+ is (κ, \mathbf{U}^+) -accessible. Using [proposition 0.2.31](#), we see that i_* is also a $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -accessible extension, so we apply the lemma above to conclude that the left Beck–Chevalley condition is satisfied. \blacksquare

Theorem 0.3.13 (Stability of accessible adjoint functors). *Let \mathbf{U} and \mathbf{U}^+ be universes, with $\mathbf{U} \in \mathbf{U}^+$, and let κ and λ be regular cardinals in \mathbf{U} , with $\kappa \leq \lambda$. Suppose:*

- \mathcal{C} is a locally κ -presentable \mathbf{U} -category.
- \mathcal{D} is a locally λ -presentable \mathbf{U} -category.
- \mathcal{C}^+ is a locally κ -presentable \mathbf{U}^+ -category.
- \mathcal{D}^+ is a locally λ -presentable \mathbf{U}^+ -category.

Let $i : \mathcal{C} \rightarrow \mathcal{C}^+$ be a $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -accessible extension and let $j : \mathcal{D} \rightarrow \mathcal{D}^+$ be a fully faithful functor.

(i) Given a strictly commutative diagram of the form below,

$$\begin{array}{ccc} \mathcal{D} & \xrightarrow{j} & \mathcal{D}^+ \\ G \downarrow & & \downarrow G^+ \\ \mathcal{C} & \xrightarrow{i} & \mathcal{C}^+ \end{array}$$

where G is (λ, \mathbf{U}) -accessible and G^+ is (λ, \mathbf{U}^+) -accessible, if both have left adjoints and j is a $(\lambda, \mathbf{U}, \mathbf{U}^+)$ -accessible extension, then the diagram satisfies the left Beck–Chevalley condition.

(ii) Given a strictly commutative diagram of the form below,

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{i} & \mathcal{C}^+ \\ F \downarrow & & \downarrow F^+ \\ \mathcal{D} & \xrightarrow{j} & \mathcal{D}^+ \end{array}$$

if both F and F^+ have right adjoints, then the diagram satisfies the right Beck–Chevalley condition.

Proof. (i). The proof is essentially the same as lemma 0.3.11, though we have to use proposition 0.3.12 to ensure that j preserves colimits for all \mathbf{U} -small κ -filtered diagrams in \mathcal{C} .

(ii). Let D be any object in \mathcal{D} . Inspecting the proof of theorem 0.2.34, we see that our hypotheses, plus the fact that i preserves colimits for all \mathbf{U} -small diagrams in \mathcal{C} , imply that the functor $(F \downarrow D) \rightarrow (F^+ \downarrow jD)$ induced by i preserves terminal objects. Thus lemma 0.3.6 implies that the diagram satisfies the right Beck–Chevalley condition. ■

Theorem 0.3.14. *If $i : \mathcal{C} \rightarrow \mathcal{C}^+$ is a $(\kappa, \mathbf{U}, \mathbf{U}^+)$ -accessible extension and \mathcal{C} is a locally κ -presentable \mathbf{U} -category, then:*

- (i) *If λ is a regular cardinal and $\kappa \leq \lambda \in \mathbf{U}$, then $i : \mathcal{C} \rightarrow \mathcal{C}^+$ is also a $(\lambda, \mathbf{U}, \mathbf{U}^+)$ -accessible extension.*
- (ii) *If μ is the cardinality of \mathbf{U} , then $i : \mathcal{C} \rightarrow \mathcal{C}^+$ factors through the inclusion $\mathbf{K}_\mu^{\mathbf{U}^+}(\mathcal{C}^+) \hookrightarrow \mathcal{C}^+$ as functor $\mathcal{C} \rightarrow \mathbf{K}_\mu^{\mathbf{U}^+}(\mathcal{C}^+)$ that is (fully faithful and) essentially surjective on objects.*
- (iii) *The (μ, \mathbf{U}^+) -accessible functor $\mathbf{Ind}_{\mathbf{U}^+}^\mu(\mathcal{C}) \rightarrow \mathcal{C}^+$ induced by $i : \mathcal{C} \rightarrow \mathcal{C}^+$ is fully faithful and essentially surjective on objects.*

Proof. (i). Since $i : \mathcal{C} \rightarrow \mathcal{C}^+$ is a (κ, \mathbf{U}) -accessible functor, it is certainly also (λ, \mathbf{U}) -accessible, by lemma 0.2.23. It is therefore enough to show that i restricts to a functor $\mathbf{K}_\kappa^{\mathbf{U}}(\mathcal{C}) \rightarrow \mathbf{K}_\kappa^{\mathbf{U}^+}(\mathcal{C}^+)$ that is (fully faithful and) essentially surjective on objects.

Proposition 0.2.30 says $\mathbf{K}_\lambda^{\mathbf{U}}(\mathcal{C})$ is the smallest replete full subcategory of \mathcal{C} that contains $\mathbf{K}_\kappa^{\mathbf{U}}(\mathcal{C})$ and is closed in \mathcal{C} under colimits for λ -small diagrams, therefore the replete closure of the image of $\mathbf{K}_\lambda^{\mathbf{U}}(\mathcal{C})$ must be the smallest replete full

subcategory of \mathcal{C}^+ that contains $\mathbf{K}_\kappa^{\mathbf{U}^+}(\mathcal{C}^+)$ and is closed in \mathcal{C}^+ under colimits for λ -small diagrams, since i is fully faithful and preserves colimits for all \mathbf{U} -small diagrams. This proves the claim.

(ii). Since every object in \mathcal{C} is (λ, \mathbf{U}) -compact for some regular cardinal $\lambda < \mu$, claim (i) implies that the image of $i : \mathcal{C} \rightarrow \mathcal{C}^+$ is contained in $\mathbf{K}_\mu^{\mathbf{U}^+}(\mathcal{C})$. To show i is essentially surjective onto $\mathbf{K}_\mu^{\mathbf{U}^+}(\mathcal{C})$, we simply have to observe that the inaccessibility of μ (proposition 0.1.35) and proposition 0.2.30 imply that, for \mathcal{C}' any (μ, \mathbf{U}^+) -compact object in \mathcal{C}^+ , there exists a regular cardinal $\lambda < \mu$ such that \mathcal{C}' is also a (λ, \mathbf{U}^+) -compact object, which reduces the question to claim (i).

(iii). This is an immediate corollary of claim (ii) and the classification theorem (0.2.19) applied to \mathcal{C}^+ , considered as a (μ, \mathbf{U}^+) -accessible category. ■

Remark 0.3.15. Although the fact $i : \mathcal{C} \rightarrow \mathcal{C}^+$ that preserves limits and colimits for all \mathbf{U} -small diagrams in \mathcal{C} is a formal consequence of the theorem above, it is not clear whether the theorem can be proven without already knowing this.

Corollary 0.3.16. *If \mathbb{B} is a \mathbf{U} -small category and has colimits for all κ -small diagrams, and μ is the cardinality of \mathbf{U} , then the canonical (μ, \mathbf{U}^+) -accessible functor $\mathbf{Ind}_{\mathbf{U}^+}^\mu(\mathbf{Ind}_{\mathbf{U}}^\kappa(\mathbb{B})) \rightarrow \mathbf{Ind}_{\mathbf{U}^+}^\kappa(\mathbb{B})$ is fully faithful and essentially surjective on objects.* ■

SIMPLICIAL SETS

Simplicial sets, like simplicial complexes, are combinatorial models for spaces built up by gluing standard n -simplices together; unlike simplicial complexes, an n -simplex in a simplicial set need not be uniquely determined by its vertices. It is for this reason that simplicial sets were once known by the unwieldy name ‘complete semi-simplicial (c.s.s.) complex’.

In the 1960s, it was discovered that one can mimic the definitions and constructions of classical homotopy theory by combinatorial means using simplicial sets, and that the resulting theory is moreover equivalent to the classical theory in a natural, functorial way. More recently, it has been shown that the homotopy theory of simplicial sets is *universal* in a precise sense,^[1] so it seems fitting that we begin here.

I.1 Basics

Definition I.1.1. The **simplex category** is the category $\mathbf{\Delta}$ whose objects are the positive finite ordinals and whose morphisms are the monotone maps. We use the geometer’s convention: $[n]$ denotes the ordinal $\{0, 1, \dots, n\}$.

Definition I.1.2. A **simplicial object** in a category \mathcal{C} is a functor $\mathbf{\Delta}^{\text{op}} \rightarrow \mathcal{C}$, and a **morphism of simplicial objects** in \mathcal{C} is a natural transformation of such functors. The **category of simplicial objects** in \mathcal{C} is the functor category $[\mathbf{\Delta}^{\text{op}}, \mathcal{C}]$ and is denoted by $\text{s}\mathcal{C}$.

^[1] See [Dugger, 2001].

Definition 1.1.3. The **coface maps** in $\mathbf{\Delta}$ are the morphisms $\delta_n^i : [n-1] \rightarrow [n]$, where δ_n^i is the unique injective monotone map that misses i ; and the **codegeneracy maps** in $\mathbf{\Delta}$ are the morphisms $\sigma_n^i : [n+1] \rightarrow [n]$, where σ_n^i is the unique surjective monotone map with $\sigma_n^i(i) = \sigma_n^i(i+1) = i$.

Theorem 1.1.4 (Cosimplicial identities). *The following equations hold in $\mathbf{\Delta}$:*

$$\begin{aligned}
 \delta_{n+1}^{j+1} \circ \delta_n^i &= \delta_{n+1}^i \circ \delta_n^j && \text{if } 0 \leq i \leq j \leq n \\
 \sigma_n^j \circ \sigma_{n+1}^i &= \sigma_n^i \circ \sigma_{n+1}^{j+1} && \text{if } 0 \leq i \leq j \leq n \\
 \sigma_{n+1}^{j+1} \circ \delta_{n+1}^i &= \delta_n^i \circ \sigma_n^j && \text{if } 0 \leq i \leq j \leq n \\
 \delta_n^{j+1} \circ \sigma_n^i &= \sigma_{n+1}^i \circ \delta_{n+1}^{j+2} && \text{if } 0 \leq i < j < n \\
 \sigma_n^i \circ \delta_n^i &= \text{id} && \text{if } 0 \leq i \leq n \\
 \sigma_n^{i+1} \circ \delta_n^i &= \text{id} && \text{if } 0 \leq i < n
 \end{aligned}$$

Equivalently, the following diagrams commute:

$$\begin{array}{ccc}
 [n-1] & \xrightarrow{\delta^i} & [n] \\
 \delta^j \downarrow & & \downarrow \delta^{j+1} \\
 [n] & \xrightarrow{\delta^i} & [n+1]
 \end{array} \quad \text{for } 0 \leq i \leq j \leq n$$

$$\begin{array}{ccc}
 [n+1] & \xrightarrow{\sigma^i} & [n] \\
 \sigma^{j+1} \downarrow & & \downarrow \sigma^j \\
 [n] & \xrightarrow{\sigma^i} & [n-1]
 \end{array} \quad \text{for } 0 \leq i \leq j \leq n$$

$$\begin{array}{ccc}
 [n] & \xrightarrow{\delta^i} & [n+1] \\
 \sigma^j \downarrow & & \downarrow \sigma^{j+1} \\
 [n-1] & \xrightarrow{\delta^i} & [n]
 \end{array} \quad \text{for } 0 \leq i \leq j \leq n$$

$$\begin{array}{ccc}
 [n] & \xrightarrow{\sigma^i} & [n-1] \\
 \delta^{j+2} \downarrow & & \downarrow \delta^{j+1} \\
 [n+1] & \xrightarrow{\sigma^i} & [n]
 \end{array} \quad \text{for } 0 \leq i < j < n$$

$$\begin{array}{ccc}
 [n-1] & \xrightarrow{\delta^i} & [n] \\
 \delta^i \downarrow & \searrow \text{id} & \downarrow \sigma^{i+1} \\
 [n] & \xrightarrow{\sigma^i} & [n-1]
 \end{array} \quad \text{for } 0 \leq i \leq n$$

Moreover, every morphism $[n] \rightarrow [m]$ in $\mathbf{\Delta}$ is uniquely a composite of the form

$$\delta_m^{j_1} \circ \dots \circ \delta_k^{j_{m-k}} \circ \sigma_k^{i_{n-k}} \circ \dots \circ \sigma_n^{i_1}$$

where $k \leq \min\{n, m\}$, and

$$\begin{aligned}
 0 &\leq i_{n-k} \leq \dots \leq i_1 \leq n \\
 0 &\leq j_{m-k} \leq \dots \leq j_1 \leq m
 \end{aligned}$$

The category $\mathbf{\Delta}$ is uniquely characterised by these properties.

Proof. See [May, 1967, § 2], [GZ, Ch. II, § 2], or [Weibel, 1994, § 8.1]. \square

Definition 1.1.5. Let A be a simplicial object in a category \mathcal{C} . A **face operator** for A is a morphism of the form $A(\delta_n^i) : A([n]) \rightarrow A([n-1])$, and a **degeneracy operator** for A is a morphism of the form $A(\sigma_n^i) : A([n]) \rightarrow A([n+1])$. For brevity, we will usually write A_n instead of $A([n])$, d_n^i instead of $A(\delta_n^i)$, and s_n^i instead of $A(\sigma_n^i)$.

Corollary 1.1.6 (Simplicial identities). *The face and degeneracy operators of a simplicial object satisfy the formal duals of the equations in theorem 1.1.4.* \blacksquare

Corollary 1.1.7. *A simplicial object A is uniquely determined by the sequence of objects A_0, A_1, A_2, \dots together with the face and degeneracy operators. Conversely, any sequence of objects equipped with face and degeneracy operators satisfying the simplicial identities defined a simplicial object.* \blacksquare

Definition 1.1.8. A **simplicial set** is a simplicial object in \mathbf{Set} , and the **category of simplicial sets** is denoted by \mathbf{sSet} .

Lemma 1.1.9.

- (i) *Limits (resp. colimits) in \mathbf{sSet} are constructed degreewise: a cone (resp. cocone) in \mathbf{sSet} over a diagram is limiting (resp. colimiting) if and only if it is so in every degree.*

- (ii) A morphism of \mathbf{sSet} is monic (resp. epic) if and only if it is degreewise injective (resp. surjective).

Proof. These are standard facts about functor categories. \square

Definition I.I.10. The **standard n -simplex** in \mathbf{sSet} , denoted by Δ^n , is the representable presheaf $\Delta(-, [n])$.

Theorem I.I.11. Let $\Delta^\bullet : \mathbf{\Delta} \rightarrow \mathbf{sSet}$ be the functor $[n] \mapsto \Delta^n$.

- (i) For any simplicial set X , the map $\mathbf{sSet}(\Delta^n, X) \rightarrow X_n$ defined by $f \mapsto f_n(\text{id}_{[n]})$ is a bijection and is moreover natural in $[n]$ and X .
- (ii) \mathbf{sSet} has limits and colimits for all small diagrams, every epimorphism is effective, and for all morphisms $f : X \rightarrow Y$ in \mathbf{sSet} , the pullback functor $f^* : \mathbf{sSet}_{/Y} \rightarrow \mathbf{sSet}_{/X}$ preserves colimits.
- (iii) Δ^\bullet is dense, i.e. for any simplicial set X , the tautological cocone from the canonical diagram $(\Delta^\bullet \downarrow X) \rightarrow \mathbf{sSet}$ to X is colimiting.
- (iv) Let \mathcal{E} be a locally small category with colimits for all small diagrams. If $F : \mathbf{sSet} \rightarrow \mathcal{E}$ is a functor that preserves small colimits, then it is left adjoint to the functor $\mathcal{E} \rightarrow \mathbf{sSet}$ defined by $E \mapsto \mathcal{E}(F\Delta^\bullet, E)$.
- (v) With \mathcal{E} as above, the functor $F \mapsto F\Delta^\bullet$ from the category of colimit-preserving functors $\mathbf{sSet} \rightarrow \mathcal{E}$ to the category of all functors $\mathbf{\Delta} \rightarrow \mathcal{E}$ is fully faithful and essentially surjective on objects.

Proof. Claim (i) is just the Yoneda lemma, claim (ii) follows from the lemma above, and claims (iii)–(v) are a special case of what is discussed in [ML–M, Ch. I, § 5]. \square

¶ I.I.12. An element of X_n is often called an **n -simplex** in X ; in particular, an element of X_0 is a **vertex** of X and an element of X_1 is an **edge** of X . This is justified by statement (i) in the above theorem.

Corollary I.I.13. There exists an adjunction

$$\tau_1 \dashv \mathbf{N} : \mathbf{Cat} \rightarrow \mathbf{sSet}$$

such that $\tau_1 \Delta^n$ is just $[n]$ regarded as a preorder category, and this adjunction is unique up to unique isomorphism. The functor \mathbf{N} is moreover fully faithful and exhibits \mathbf{Cat} as a reflective subcategory of \mathbf{sSet} .

Proof. In this case it is easier to use the general adjoint functor theorem to construct a left adjoint for the functor N defined by $N(\mathbb{C})_n = \text{ob} [[n], \mathbb{C}]$. ■

Definition I.1.14. The **fundamental category** of a simplicial set X is the small category $\tau_1 X$, and the **nerve** of a small category \mathbb{C} is the simplicial set $N(\mathbb{C})$.

Remark I.1.15. Given a simplicial set X , the fundamental category $\tau_1 X$ admits the following presentation by generators and relations: the objects are the vertices of X , and the morphisms are generated by the edges of X , modulo the relation $d_0(x) \circ d_2(x) = d_1(x)$ for all 2-simplices x in X .

Corollary I.1.16. *There exists an adjunction*

$$\pi_0 \dashv \text{disc} : \mathbf{Set} \rightarrow \mathbf{sSet}$$

such that $\pi_0 \Delta^n = 1$ for all n , and this adjunction is unique up to unique isomorphism. Explicitly, we may take

$$(\text{disc } Y)_n = Y$$

for all n , with id_Y for all the face and degeneracy operators. The functor disc is moreover fully faithful and exhibits \mathbf{Set} as a reflective subcategory of \mathbf{sSet} . ■

Definition I.1.17. The **set of connected components** of a simplicial set X is the set $\pi_0 X$, and a **discrete simplicial set** is one that is isomorphic to $\text{disc } Y$ for some set Y .

¶ **I.1.18.** We will usually not distinguish between Y and $\text{disc } Y$ notationally.

Definition I.1.19. The **standard n -simplex** in \mathbf{Top} , denoted by $|\Delta^n|$, is the topological space

$$|\Delta^n| = \{(x_0, \dots, x_n) \in [0, 1]^{n+1} \mid x_0 + \dots + x_n = 1\}$$

where $[0, 1]$ is the closed unit interval with the standard metric. The functor $|\Delta^\bullet| : \mathbf{\Delta} \rightarrow \mathbf{Top}$ sends $[n]$ to $|\Delta^n|$ and is defined on morphisms by linearly interpolating the obvious map of vertices.

Corollary I.1.20. *There exists an adjunction*

$$|_|\dashv S : \mathbf{Top} \rightarrow \mathbf{sSet}$$

extending the functor $|\Delta^\bullet| : \mathbf{\Delta} \rightarrow \mathbf{Top}$ defined above, and this adjunction is unique up to unique isomorphism. ■

Definition 1.1.21. The **geometric realisation** of a simplicial set X is the topological space $|X|$, and the **singular set** of a topological space Y is the simplicial set $S(Y)$.

Definition 1.1.22. Let \mathcal{C} be a category with binary products, and let Y and Z be objects in \mathcal{C} . An **exponential object** for Y and Z is an object Z^Y in \mathcal{C} and a morphism $\text{ev}_{Y,Z} : Z^Y \times Y \rightarrow Z$ with the following universal property:

- For all morphisms $f : X \times Y \rightarrow Z$ in \mathcal{C} , there exists a unique morphism $\bar{f} : X \rightarrow Z^Y$ such that $\text{ev}_{Y,Z} \circ (\bar{f} \times \text{id}_Y) = f$.

A **cartesian closed category** is a category with all finite products and exponential objects for all pairs of objects.

Proposition 1.1.23. Let **CGHaus** be the category of compactly-generated Hausdorff spaces^[1] and continuous maps.

- (i) **sSet** is a cartesian closed category, with exponential objects given by the formula below:

$$Z^Y = \mathbf{sSet}(\Delta^\bullet \times Y, Z)$$

- (ii) If Y is a locally compact Hausdorff space, then for all topological spaces Z , the set of all continuous maps $Y \rightarrow Z$, equipped with the compact-open topology, is an exponential object Z^Y in **Top**.
- (iii) **CGHaus** is a cartesian closed category.

Proof. Claim (i) can be verified by direct calculation, claim (ii) follows from Theorems 46.10 and 46.11 in [Munkres, 2000], and claim (iii) is proved in [GZ, Ch. III, § 2]. □

Theorem 1.1.24.

- (i) The topological standard n -simplex $|\Delta^n|$ is a compact Hausdorff space.
- (ii) For any simplicial set X , the geometric realisation $|X|$ is a compactly-generated Hausdorff space.
- (iii) The previously-constructed adjunction $|-| \dashv S : \mathbf{Top} \rightarrow \mathbf{sSet}$ restricts to an adjunction between **CGHaus** and **sSet**, and moreover the functor $|-| : \mathbf{sSet} \rightarrow \mathbf{CGHaus}$ preserves finite limits and reflects isomorphisms.

^[1] — also known as **Kelley spaces**.

Proof. Claim (i) is a standard fact, while claims (ii) and (iii) are proven in [GZ, Ch. III, § 3]. □

1.2 The Quillen model structure

Prerequisites. §§ 1.1, A.1.

In [1967], Quillen constructed an axiomatic framework for doing homotopy theory in abstract categories, which he called ‘closed model categories’, and showed that \mathbf{sSet} can be endowed with a model structure such that the resulting homotopy theory is equivalent in a strong sense to the homotopy theory of topological spaces.

Definition 1.2.1. A **horn** is a simplicial subset of the form $\Lambda_k^n \subseteq \Delta^n$, where Λ_k^n is the union of the images of $\delta_n^0, \dots, \delta_n^{k-1}, \delta_n^{k+1}, \dots, \delta_n^n : \Delta^{n-1} \rightarrow \Delta^n$ in \mathbf{sSet} . In other words, Λ_k^n is the union of all the faces of Δ^n that include the k -th vertex. The **boundary** of Δ^n is the simplicial subset $\partial\Delta^n \subseteq \Delta^n$ generated by the images of $\delta_n^0, \dots, \delta_n^n : \Delta^{n-1} \rightarrow \Delta^n$.

Definition 1.2.2. A **cofibration** in \mathbf{sSet} is a monomorphism. A **Kan fibration** is a morphism $f : X \rightarrow Y$ in \mathbf{sSet} that has the right lifting property with respect to the horn inclusions $\Lambda_k^n \hookrightarrow \Delta^n$, where $n \geq 1$ and $0 \leq k \leq n$. A **Kan complex** is a simplicial set X such that the unique morphism $X \rightarrow 1$ is a Kan fibration.

Proposition 1.2.3. *Let C_Q and F_Q be, respectively, the class of cofibrations and the class of Kan fibrations in \mathbf{sSet} . There exists a unique class \mathcal{W}_Q of morphisms in \mathbf{sSet} with the following properties:*

- (i) $(C_Q \cap \mathcal{W}_Q, F_Q)$ is a weak factorisation system for \mathbf{sSet} .
- (ii) $(C_Q, F_Q \cap \mathcal{W}_Q)$ is a weak factorisation system for \mathbf{sSet} .
- (iii) A morphism is in \mathcal{W}_Q if and only if it is of the form $f \circ c$, where f is in F_Q and has the right lifting property with respect to all morphisms in C_Q , and c is in C_Q and has the left lifting property with respect to all morphisms in F_Q .

Proof. ??? □

Definition 1.2.4. A **Quillen weak equivalence** in \mathbf{sSet} is a morphism in the class \mathcal{W}_Q described above. An **anodyne extension**, or **trivial cofibration** in \mathbf{sSet} , is a cofibration that has the left lifting property with respect to all Kan fibrations. A **trivial Kan fibration** is a Kan fibration that has the right lifting property with respect to all cofibrations.

HOMOTOPICAL CATEGORIES

2.1 Basics

Prerequisites. § A.2.

Definition 2.1.1. A relative category \mathcal{C} is a **category with weak equivalences** if $\text{weq } \mathcal{C}$ has the 2-out-of-3 property, and it is a **homotopical category** if $\text{weq } \mathcal{C}$ has the 2-out-of-6 property. A **homotopical functor** is a relative functor between homotopical categories.

Example 2.1.2. Any saturated relative category is automatically a homotopical category, by [corollary A.2.14](#). In particular, any minimal saturated relative category is a homotopical category. On the other hand, any maximal relative category is obviously a homotopical category.

Remark 2.1.3. A relative category \mathcal{C} is a category with weak equivalences or a homotopical category if and only if the opposite relative category \mathcal{C}^{op} is.

Lemma 2.1.4. *Let A be an object in a homotopical category (resp. category with weak equivalences) \mathcal{C} . Then the slice category $\mathcal{C}_{/A}$ is also a homotopical category (resp. category with weak equivalences) if we declare a morphism in $\mathcal{C}_{/A}$ to be a weak equivalence if and only if it is a weak equivalence in \mathcal{C} .*

Proof. Use [lemma A.2.13](#) on the projection functor $\mathcal{C}_{/A} \rightarrow \mathcal{C}$. ■

Definition 2.1.5. Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be two (not necessarily relative) functors between relative categories. A **natural weak equivalence** $\alpha : F \Rightarrow G$ is a natural transformation such that $\alpha_{\mathcal{C}} : FC \rightarrow GC$ is a weak equivalence in \mathcal{D} for

all objects C in \mathcal{C} , and we say F and G are **naturally weakly equivalent** if they can be connected by a *zigzag* of natural weak equivalences.

Remark 2.1.6. If F and G are relative functors, then this is precisely the notion of weak equivalence in the relative functor category $[\mathcal{C}, \mathcal{D}]_h$. Although the definition above applies to all functors, if $H : \mathcal{D} \rightarrow \mathcal{E}$ is a functor, then the natural transformation $H\alpha : HF \Rightarrow HG$ is only guaranteed to be a natural weak equivalence if we assume H is a relative functor.

Definition 2.1.7. A **homotopical equivalence** is a relative functor $F : \mathcal{C} \rightarrow \mathcal{D}$ for which there exists a relative functor $G : \mathcal{D} \rightarrow \mathcal{C}$ such that GF is naturally weakly equivalent to $\text{id}_{\mathcal{C}}$ and FG is naturally weakly equivalent to $\text{id}_{\mathcal{D}}$. Such a G is said to be a **homotopical inverse** of F .

Proposition 2.1.8. *If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a homotopical equivalence of relative categories with homotopical inverse $G : \mathcal{D} \rightarrow \mathcal{C}$, then $\text{Ho } F : \text{Ho } \mathcal{C} \rightarrow \text{Ho } \mathcal{D}$ is an equivalence of categories, with quasi-inverse $\text{Ho } G : \text{Ho } \mathcal{D} \rightarrow \text{Ho } \mathcal{C}$. ■*

2.2 Homotopical Kan extensions

Prerequisites. § 2.1.

Definition 2.2.1. Let \mathcal{C} be a homotopical category. A **homotopically initial object** in \mathcal{C} is an object A for which there exists a zigzag of natural transformations of the form

$$\Delta A \text{ ——— } F \xrightarrow{\alpha} G \text{ ——— } \text{id}_{\mathcal{C}}$$

where $\Delta A : \mathcal{C} \rightarrow \mathcal{C}$ is the constant functor with value A , $\alpha_A : FA \rightarrow GA$ is a weak equivalence in \mathcal{C} , and the unmarked lines denote (possibly trivial) zigzags of natural weak equivalences. Dually, a **homotopically terminal object** in \mathcal{C} is a homotopically initial object in \mathcal{C}^{op} .

Proposition 2.2.2. *Let \mathcal{C} be a homotopical category. If A is a homotopically initial (resp. homotopically terminal) object in \mathcal{C} , then:*

- (i) *Any object in \mathcal{C} weakly equivalent to A is also a homotopically initial (resp. homotopically terminal) object in \mathcal{C} .*
- (ii) *A is an initial (resp. terminal) object in $\text{Ho } \mathcal{C}$.*

(iii) *If C is a minimal homotopical category, then A is an initial (resp. terminal) object in C as well.*

Conversely, any initial (resp. terminal) object in C is also homotopically initial (resp. homotopically terminal).

Proof. Obvious. (This is Proposition 38.3 in [DHKS].) ■

Definition 2.2.3. A **homotopically contractible category** is a homotopical category C such that the unique (homotopical) functor $C \rightarrow \mathbb{1}$ is a homotopical equivalence, where $\mathbb{1}$ is the trivial category with only one object.

Proposition 2.2.4. *Let C be a homotopical category. The following are equivalent:*

- (i) *C is homotopically contractible.*
- (ii) *C is inhabited, and for every object A in C , the constant functor ΔA is naturally weakly equivalent to id_C .*
- (iii) *There exists an object A in C such that ΔA and id_C are naturally weakly equivalent.*

Proof. Obvious. (This is paragraph 37.6 in [DHKS].) ■

Proposition 2.2.5. *Let C be a homotopically contractible category.*

- (i) *Every morphism in C is a weak equivalence.*
- (ii) *The unique functor $\text{Ho } C \rightarrow \mathbb{1}$ is an equivalence of categories.*
- (iii) *If C is a minimal homotopical category, then $C \rightarrow \mathbb{1}$ is also an equivalence of categories.*
- (iv) *The opposite homotopical category C^{op} and the homotopical functor category $[\mathcal{D}, C]_{\text{h}}$ (for any homotopical category \mathcal{D}) are also homotopically contractible.*
- (v) *Every object in C is both homotopically initial and homotopically terminal.*

Proof. Obvious. (This is paragraph 37.6 in [DHKS].) ■

Proposition 2.2.6. *Let \mathcal{C} be a homotopical category. If \mathcal{D} is the full homotopical subcategory of \mathcal{C} spanned by the homotopically initial (or homotopically terminal) objects, then \mathcal{D} is homotopically contractible.*

Proof. See paragraph 38.5 in [DHKS]. □

Definition 2.2.7. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{C} \rightarrow \mathcal{E}$ be two (not necessarily homotopical) functors between homotopical categories. A **homotopical left Kan extension** (resp. **homotopical right Kan extension**) of G along F is a homotopically initial (resp. homotopically terminal) object of the relative category $(G \downarrow F^*)_h$ (resp. $(F^* \downarrow G)_h$) described below:

- The objects are pairs (H, α) where H is a homotopical functor $\mathcal{D} \rightarrow \mathcal{E}$ and α is a natural transformation of type $G \Rightarrow HF$ (resp. $HF \Rightarrow G$).
- The morphisms $(H', \alpha') \rightarrow (H, \alpha)$ are those natural transformations $\beta : H' \Rightarrow H$ such that $\beta F \bullet \alpha' = \alpha$ (resp. $\alpha \bullet \beta F = \alpha'$).
- The weak equivalences are the natural weak equivalences.

MODEL CATEGORIES

3.1 Basics

Prerequisites. §§ 2.1, A.1.

In [1967], Quillen introduced the notion of a ‘closed model category’ (but we shall say simply ‘model category’) for homotopy theory, so as to formalise the similarities between the homotopy theory of spaces and homological algebra. The idea was that, to do homotopy theory, one really only needed to know which morphisms are cofibrations, which are weak equivalences, and which are fibrations.

Definition 3.1.1. A **model category** is a locally small category \mathcal{M} equipped with three subclasses $\mathcal{C}, \mathcal{W}, \mathcal{F}$ of $\text{mor } \mathcal{M}$ satisfying the following axioms:

- **CM1.** \mathcal{M} has finite limits and finite colimits.
- **CM2.** \mathcal{W} has the 2-out-of-3 property.
- **CM3.** $\mathcal{C}, \mathcal{W},$ and \mathcal{F} are closed under retracts.
- **CM4.** Given a commutative diagram

$$\begin{array}{ccc} A & \longrightarrow & X \\ i \downarrow & & \downarrow p \\ B & \longrightarrow & Y \end{array}$$

where i is in \mathcal{C} and p is in \mathcal{F} , if at least one of i or p is also in \mathcal{W} , then there exists a morphism $B \rightarrow X$ making the evident triangles commute.

- **CM5.** Any morphism f in \mathcal{M} may be factored in two ways:
 - $f = p \circ i$ where i is in $C \cap \mathcal{W}$ and p is in \mathcal{F} , and
 - $f = q \circ j$, where j is in C and q is in $\mathcal{W} \cap \mathcal{F}$.

The triple $(C, \mathcal{W}, \mathcal{F})$ is said to be a **model structure** on \mathcal{M} . Given such a model structure on \mathcal{M} ,

- a **cofibration** is a morphism in C ,
- a **weak equivalence** is a morphism in \mathcal{W} ,
- a **fibration** is a morphism in \mathcal{F} ,
- a **trivial cofibration** (or **acyclic cofibration**) is a morphism in $C \cap \mathcal{W}$, and
- a **trivial fibration** (or **acyclic fibration**) is a morphism in $\mathcal{W} \cap \mathcal{F}$;
- a **cofibrant object** in \mathcal{M} is an object X such that the unique morphism $0 \rightarrow X$ is a cofibration, and
- a **fibrant object** in \mathcal{M} is an object X such that the unique morphism $X \rightarrow 1$ is a fibration.
- a **cofibrant–fibrant object** in \mathcal{M} is an object that is both cofibrant and fibrant.

Remark 3.1.2. The above presentation of the axioms is due to Quillen [1969], and is the one used in [DS] and [GJ]; however, [DHKS], [Hirschhorn, 2003], and [Hovey, 1999] use a variant definition that replaces axioms CM1 and CM5 with stronger ones:

- **CM1'**. \mathcal{M} is complete and cocomplete.
- **CM5'**. The $(C \cap \mathcal{W}, \mathcal{F})$ and $(C, \mathcal{W} \cap \mathcal{F})$ -factorisations can be chosen *functorially* in the sense of [definition A.1.9](#).

Note also that Hovey [1999] considers the functorial factorisations to be a *structure* rather than a property.

Remark 3.1.3. Let \mathcal{M} be a category with finite limits and finite colimits. Then, $(C, \mathcal{W}, \mathcal{F})$ is a model structure on \mathcal{M} if and only if $(\mathcal{F}^{\text{op}}, \mathcal{W}^{\text{op}}, C^{\text{op}})$ is a model structure on \mathcal{M}^{op} .

Theorem 3.1.4. *Let \mathcal{M} be a locally small category and let $\mathcal{C}, \mathcal{W}, \mathcal{F}$ be subclasses of $\text{mor } \mathcal{M}$. Assuming \mathcal{M} has finite limits and finite colimits, the following are equivalent:*

- (i) $(\mathcal{C}, \mathcal{W}, \mathcal{F})$ is a model structure for \mathcal{M} .
- (ii) \mathcal{M} is a saturated homotopical category with $\text{weq } \mathcal{M} = \mathcal{W}$, and both $(\mathcal{C} \cap \mathcal{W}, \mathcal{F})$ and $(\mathcal{C}, \mathcal{W} \cap \mathcal{F})$ are weak factorisation systems for \mathcal{M} .
- (iii) $(\mathcal{M}, \mathcal{W})$ is a category with weak equivalences (as in [definition 2.1.1](#)), and both $(\mathcal{C} \cap \mathcal{W}, \mathcal{F})$ and $(\mathcal{C}, \mathcal{W} \cap \mathcal{F})$ are weak factorisation systems for \mathcal{M} .

Proof. (i) \Rightarrow (ii). The fact that we have two weak factorisation systems follows from Lemma 1.1 in [GJ, Ch. II] or Proposition 7.2.3 in [Hirschhorn, 2003]; and the saturation property follows from Theorems 1.10 and 1.11 in [GJ, Ch. II], or Theorem 8.3.10 in [Hirschhorn, 2003].

(ii) \Rightarrow (iii). Obvious.

(iii) \Rightarrow (i). Use [proposition A.1.7](#). □

Lemma 3.1.5. *Let A be an object in a model category \mathcal{M} . Then the slice category $\mathcal{M}_{/A}$ has the **slice model structure**, where a morphism in $\mathcal{M}_{/A}$ is a cofibration, weak equivalence, or fibration if it is so in \mathcal{M} .*

Proof. Use lemmas [2.1.4](#) and [A.1.6](#), plus the fact that $\mathcal{M}_{/A}$ has finite limits and finite colimits if \mathcal{M} does. ■

Definition 3.1.6. A **left Quillen functor** is a functor $F : \mathcal{M} \rightarrow \mathcal{N}$ between model categories that has a right adjoint and preserves cofibrations and trivial cofibrations; dually, a **right Quillen functor** is a functor $G : \mathcal{N} \rightarrow \mathcal{M}$ between model categories that has a left adjoint and preserves fibrations and trivial fibrations. A **Quillen adjunction** is an adjunction

$$F \dashv G : \mathcal{M} \rightarrow \mathcal{N}$$

where \mathcal{M} and \mathcal{N} are model categories, such that F is a left Quillen functor and G is a right Quillen functor. A **Quillen equivalence** is a Quillen adjunction as above satisfying this additional condition:

- Given a cofibrant object A in \mathcal{N} and fibrant object X in \mathcal{M} , a morphism $FA \rightarrow X$ is a weak equivalence in \mathcal{M} if and only if its adjoint transpose $A \rightarrow GX$ is a weak equivalence in \mathcal{N} .

Proposition 3.1.7. *Let $F \dashv G : \mathcal{M} \rightarrow \mathcal{N}$ be an adjunction between model categories. The following are equivalent:*

- (i) $F \dashv G$ is a Quillen adjunction.
- (ii) F is a left Quillen functor.
- (iii) G is a right Quillen functor.
- (iv) F preserves cofibrations and G preserves fibrations.
- (v) F preserves trivial cofibrations and G preserves trivial fibrations.

Proof. Use [proposition A.1.8](#). ■

Lemma 3.1.8 (Kenneth S. Brown). *Let \mathcal{M} be a model category and let \mathcal{C} be a category with weak equivalences. If $F : \mathcal{M} \rightarrow \mathcal{C}$ sends trivial cofibrations (resp. trivial fibrations) in \mathcal{M} to weak equivalences in \mathcal{C} , then F preserves all weak equivalences between cofibrant (resp. fibrant) objects.*

Proof. See Lemma 9.9 in [DS], Lemma 7.7.1 in [Hirschhorn, 2003], or Lemma 14.5 in [DHKS]. □

Corollary 3.1.9. *Let $F \dashv G : \mathcal{M} \rightarrow \mathcal{N}$ be a Quillen adjunction.*

- (i) *If A and B are cofibrant objects in \mathcal{N} and $f : A \rightarrow B$ is a weak equivalence in \mathcal{N} , then Ff is a weak equivalence in \mathcal{M} .*
- (ii) *If X and Y are fibrant objects in \mathcal{M} and $g : X \rightarrow Y$ is a weak equivalence in \mathcal{M} , then Gg is a weak equivalence in \mathcal{N} .* ■

Proposition 3.1.10 (Dugger). *Let $F \dashv G$ be an adjunction between [strong???] model categories. The following are equivalent:*

- (i) $F \dashv G$ is a Quillen adjunction.
- (ii) F preserves cofibrations between cofibrant objects and all trivial cofibrations.

(iii) G preserves fibrations between fibrant objects and all trivial fibrations.

Proof. This is Proposition 8.5.4 in [Hirschhorn, 2003]. □

Definition 3.1.11. Let X be an object in a model category \mathcal{M} .

- A **cofibrant replacement** for X is a pair (\tilde{X}, p) where \tilde{X} is a cofibrant object in \mathcal{M} and p is a weak equivalence $\tilde{X} \rightarrow X$.
- A **fibrant replacement** for X is a pair (\hat{X}, i) where \hat{X} is a fibrant object in \mathcal{M} and i is a weak equivalence $X \rightarrow \hat{X}$.
- A **fibrant cofibrant replacement** for X is a cofibrant replacement (\tilde{X}, p) where $p : \tilde{X} \rightarrow X$ is a trivial fibration.
- A **cofibrant fibrant replacement** for X is a fibrant replacement (\hat{X}, i) where $i : X \rightarrow \hat{X}$ is a trivial cofibration.

Remark 3.1.12. Note that a fibrant cofibrant replacement for X is precisely a cofibrant replacement for X that is fibrant as an object in $\mathcal{M}_{/X}$, and a cofibrant fibrant replacement for X is precisely a fibrant replacement for X that is cofibrant as an object in ${}^X/\mathcal{M}$.

Moreover, if X is fibrant and (\tilde{X}, p) is a fibrant cofibrant replacement for X , then \tilde{X} is both fibrant and cofibrant in \mathcal{M} , and if X is cofibrant and (\hat{X}, i) is a cofibrant fibrant replacement for X , then \hat{X} is both cofibrant and fibrant in \mathcal{M} .

Proposition 3.1.13. *Any object in a model category has both a fibrant cofibrant replacement and a cofibrant fibrant replacement.*

Proof. Use axiom CM5. ■

3.2 Left and right homotopy

Prerequisites. § 3.1.

Definition 3.2.1. Let X be an object in a model category \mathcal{M} . A **cylinder object** for X is a quadruple $(\text{Cyl}(X), i_0, i_1, p)$, where $\text{Cyl}(X)$ is an object in \mathcal{M} , $p : \text{Cyl}(X) \rightarrow X$ is a weak equivalence, and $i_0, i_1 : X \rightarrow \text{Cyl}(X)$ are sections of p such that the morphism $[i_0, i_1] : X + X \rightarrow \text{Cyl}(X)$ is a cofibration. Dually, a **path object** for X is a quadruple $(\text{Path}(X), i, p_0, p_1)$, where $\text{Path}(X)$ is an object

in \mathcal{M} , $i : X \rightarrow \text{Path}(X)$ is a weak equivalence, and $p_0, p_1 : \text{Path}(X) \rightarrow X$ are retractions of i such that the morphism $\langle p_0, p_1 \rangle : \text{Path}(X) \rightarrow X \times X$ is a fibration.

Proposition 3.2.2. *Let X be an object in a model category \mathcal{M} .*

- (i) *There exists a cylinder object $(\text{Cyl}(X), i_0, i_1, p)$ for X , where the morphism $p : \text{Cyl}(X) \rightarrow X$ is a trivial fibration.*
- (ii) *There exists a path object $(\text{Path}(X), i, p_0, p_1)$ for X , where the morphism $i : X \rightarrow \text{Path}(X)$ is a trivial cofibration.*

Proof. Use axioms CM1 and CM5. ■

Definition 3.2.3. Let $f_0, f_1 : X \rightarrow Y$ be a parallel pair of morphisms in a model category \mathcal{M} . A **left homotopy** from f_0 to f_1 with respect to a cylinder object $(\text{Cyl}(X), i_0, i_1, p)$ is a morphism $H : \text{Cyl}(X) \rightarrow Y$ such that $H \circ i_0 = f_0$ and $H \circ i_1 = f_1$. Dually, a **right homotopy** from f_0 to f_1 with respect to a path object $(\text{Path}(Y), i, p_0, p_1)$ is a morphism $H : X \rightarrow \text{Path}(Y)$ such that $p_0 \circ H = f_0$ and $p_1 \circ H = f_1$. We say f_0 and f_1 are **left homotopic** if there exists a left homotopy from f_0 to f_1 with respect to some cylinder object for X , and we say f_0 and f_1 are **right homotopic** if there exists a right homotopy from f_0 to f_1 with respect to some path object for Y .

Remark 3.2.4. If f_0 and f_1 are either left homotopic or right homotopic, then they must represent the same morphism in $\text{Ho } \mathcal{M}$. For definiteness, let us write $\gamma : \mathcal{M} \rightarrow \text{Ho } \mathcal{M}$ for the universal functor, and suppose $H : \text{Cyl}(X) \rightarrow Y$ is a left homotopy from f_0 to f_1 . Since i_0 and i_1 are both sections of the weak equivalence $p : \text{Cyl}(X) \rightarrow X$, we must have $\gamma i_0 = (\gamma p)^{-1} = \gamma i_1$; but $f_0 = H \circ i_0$ and $f_1 = H \circ i_1$, so indeed $\gamma f_0 = \gamma f_1$. This is one of the reasons for calling $\text{Ho } \mathcal{M}$ the *homotopy category* of \mathcal{M} .

However, it is not quite true that $\gamma f_0 = \gamma f_1$ if and only if f_0 and f_1 are either left homotopic or right homotopic; this only happens in special cases. In general, being left/right homotopic fails to even be an equivalence relation.


Lemma 3.2.5. *Let $f_0, f_1 : X \rightarrow Y$ be a parallel pair of morphisms in a model category \mathcal{M} .*

- (i) *Given any cylinder object $(\text{Cyl}(X), i_0, i_1, p)$ for X , $f_0 \circ p : \text{Cyl}(X) \rightarrow Y$ is a left homotopy from f_0 to itself.*

- (ii) Given any path object $(\text{Path}(Y), i, p_0, p_1)$ for Y , $i \circ f_0 : X \rightarrow \text{Path}(Y)$ is a right homotopy from f_0 to itself.
- (iii) If $H : \text{Cyl}(X) \rightarrow Y$ is a left homotopy from f_0 to f_1 with respect to a cylinder object $(\text{Cyl}(X), i_0, i_1, p)$ for X , then the same H is a left homotopy from f_1 to f_0 for the cylinder object $(\text{Cyl}(X), i_1, i_0, p)$.
- (iv) If $H : X \rightarrow \text{Path}(Y)$ is a right homotopy from f_0 to f_1 with respect to a path object $(\text{Path}(Y), i, p_0, p_1)$ for Y , then the same H is a right homotopy from f_1 to f_0 for the path object $(\text{Path}(Y), i, p_1, p_0)$.


Proof. Obvious. ■

Lemma 3.2.6. *Let X be a cofibrant object in a model category \mathcal{M} . Given two cylinder objects for X , say $(\text{Cyl}(X)', i'_0, i'_1, p')$ and $(\text{Cyl}(X)'', i''_0, i''_1, p'')$, there exists a third cylinder object $(\text{Cyl}(X), i_0, i_1, p)$ such that the diagram below commutes,*



and the diamond is a pushout diagram.

Dually, if Y is a fibrant object in \mathcal{M} , and we have two path objects for Y , say $(\text{Path}(Y)', i', p'_0, p'_1)$ and $(\text{Path}(Y)'', i'', p''_0, p''_1)$, then there exists a third path object $(\text{Path}(Y), i, p_0, p_1)$ such that the diagram below commutes,



and the diamond is a pullback diagram.

Proof. See Lemma 1.5 in [GJ, Ch. II], or Lemma 7.4.2 in [Hirschhorn, 2003]. □

Corollary 3.2.7. *Let $f_0, f_1, f_2 : X \rightarrow Y$ be three parallel morphisms in a model category \mathcal{M} .*

- (i) *If f_0 and f_1 are left homotopic, and f_1 and f_2 are left homotopic, then f_0 and f_2 are also left homotopic.*
- (ii) *If f_0 and f_1 are right homotopic, and f_1 and f_2 are right homotopic, then f_0 and f_2 are also right homotopic. ■*

Lemma 3.2.8. *Let $f_0, f_1 : X \rightarrow Y$ be a parallel pair of morphisms in a model category \mathcal{M} .*

- (i) *If X is cofibrant, and f_0 and f_1 are left homotopic, given any path object $(\text{Path}(Y), i, p_0, p_1)$ for Y , there is a right homotopy $H : X \rightarrow \text{Path}(Y)$ from f_0 to f_1 .*
- (ii) *If Y is fibrant, and f_0 and f_1 are right homotopic, given any cylinder object $(\text{Cyl}(X), i_0, i_1, p)$ for X , there is a left homotopy $H : \text{Cyl}(X) \rightarrow Y$ from f_0 to f_1 .*

Proof. See Proposition 1.8 in [GJ, Ch. II], or Proposition 7.4.7 in [Hirschhorn, 2003]. □

Proposition 3.2.9. *Let X and Y be objects in a model category \mathcal{M} .*

- (i) *If X is cofibrant, then being left homotopic is an equivalence relation on the hom-set $\mathcal{M}(X, Y)$.*
- (ii) *If Y is fibrant, then being right homotopic is an equivalence relation on the hom-set $\mathcal{M}(X, Y)$.*
- (iii) *If X is cofibrant and Y is fibrant, then these two equivalence relations on $\mathcal{M}(X, Y)$ coincide.*

Proof. Use the preceding lemmas. ■

Lemma 3.2.10. *Let $f_0, f_1 : X \rightarrow Y$ be a parallel pair of morphisms in a model category \mathcal{M} .*

- (i) If f_0 and f_1 are right homotopic and $g : W \rightarrow X$ is any morphism in \mathcal{M} , then $f_0 \circ g$ and $f_1 \circ g$ are also right homotopic.
- (ii) If f_0 and f_1 are left homotopic and $g : Y \rightarrow Z$ is any morphism in \mathcal{M} , then $g \circ f_0$ and $g \circ f_1$ are also left homotopic.

Proof. Obvious. ■

Corollary 3.2.11. *Let \mathcal{M} be a model category, and let \mathcal{M}_{cf} be the full subcategory spanned by the cofibrant–fibrant objects. Then the equivalence relation induced by homotopy is a congruence on \mathcal{M}_{cf} ; in particular, there exist a locally small category \mathcal{M}' and a full functor $\mathcal{M}_{\text{cf}} \rightarrow \mathcal{M}'$ with these properties:*

- The objects of \mathcal{M}' are those of \mathcal{M}_{cf} .
- The hom-set $\mathcal{M}'(X, Y)$ is $\mathcal{M}(X, Y)$ modulo homotopy.
- The functor $\mathcal{M}_{\text{cf}} \rightarrow \mathcal{M}'$ sends each morphism in \mathcal{M}' to its homotopy class. ■

The next result is a version of Whitehead’s theorem; however, this is a purely formal consequence of the model category axioms and has no real content, unlike the original theorem.

Proposition 3.2.12. *Let X and Y be cofibrant–fibrant objects in a model category \mathcal{M} . If $f : X \rightarrow Y$ is a weak equivalence, then f has a **homotopy inverse** in \mathcal{M} , i.e. a morphism $g : Y \rightarrow X$ such that $g \circ f$ and id_X are homotopic, and $f \circ g$ and id_Y are homotopic.*

Proof. See Theorem 1.10 in [GJ, Ch. II], or Theorem 7.5.10 in [Hirschhorn, 2003]. □

Corollary 3.2.13. *Let W, X, Y, Z be cofibrant–fibrant objects in a model category \mathcal{M} , and let $f_0, f_1 : X \rightarrow Y$ be a parallel pair of morphisms.*

- (i) *If $g : W \rightarrow X$ is a weak equivalence such that $f_0 \circ g$ and $f_1 \circ g$ are homotopic, then f_0 and f_1 are homotopic.*
- (ii) *If $g : Y \rightarrow Z$ is a weak equivalence such that $g \circ f_0$ and $g \circ f_1$ are homotopic, then f_0 and f_1 are homotopic.*

Proof. Use a homotopy inverse to cancel g . ■

3.3 The homotopy category

Prerequisites. §§ 3.1, 3.2, A.2.

Definition 3.3.1. The **Quillen homotopy category** (or, more simply, **homotopy category**) of a model category \mathcal{M} is the category $\text{Ho } \mathcal{M}$ obtained by freely inverting the weak equivalences in \mathcal{M} , as in [definition A.2.9](#).

Theorem 3.3.2. *Let \mathcal{M} be a model category and let $\gamma : \mathcal{M} \rightarrow \text{Ho } \mathcal{M}$ be the universal functor.*

- (i) *$\text{Ho } \mathcal{M}$ is equivalent to the locally small category \mathcal{M}' defined in [corollary 3.2.11](#), and \mathcal{M} is a saturated homotopical category.*
- (ii) *If X and Y are cofibrant–fibrant objects in \mathcal{M} , then the hom-class map $\mathcal{M}(X, Y) \rightarrow \text{Ho } \mathcal{M}(X, Y)$ induced by γ is surjective; and moreover for any parallel pair $f_0, f_1 : X \rightarrow Y$ in \mathcal{M} , we have $\gamma f_0 = \gamma f_1$ if and only if f_0 and f_1 are homotopic.*
- (iii) *For any two objects X and Y in \mathcal{M} , every morphism $X \rightarrow Y$ in $\text{Ho } \mathcal{M}$ can be represented as a zigzag of the form*

$$X \xleftarrow{p} \tilde{X} \longrightarrow \hat{Y} \xleftarrow{i} Y$$

where (\tilde{X}, p) is any cofibrant replacement for X and (\hat{Y}, i) is any fibrant replacement for Y .

Proof. (i). This is Theorem 1.11 in [GJ, Ch. II], or Proposition 5.8 in [DS].

(ii). Implied by claim (i).

(iii). Using claim (ii), every morphism $X \rightarrow Y$ in $\text{Ho } \mathcal{M}$ can be represented as a zigzag of the form

$$\begin{array}{ccc} X \xleftarrow{p} \tilde{X} & & \hat{Y} \xleftarrow{i} Y \\ & \downarrow i' & \uparrow p' \\ & R\tilde{X} \xrightarrow{f'} Q\hat{Y} & \end{array}$$

where $(R\tilde{X}, i')$ is a cofibrant fibrant replacement for \tilde{X} and $(Q\hat{Y}, p')$ is a fibrant cofibrant replacement for \hat{Y} ; but such a zigzag is manifestly equivalent to the zigzag

$$X \xleftarrow{p} \tilde{X} \xrightarrow{f} \hat{Y} \xleftarrow{i} Y$$

where $f = p' \circ f' \circ i'$. □

Corollary 3.3.3. *Let \mathcal{M} be a model category and let $\gamma : \mathcal{M} \rightarrow \text{Ho } \mathcal{M}$ be the universal functor. If X is a cofibrant object in \mathcal{M} and Y is a fibrant object in \mathcal{M} , then the hom-class map $\mathcal{M}(X, Y) \rightarrow \text{Ho } \mathcal{M}(X, Y)$ induced by γ is surjective; and moreover for any parallel pair $f_0, f_1 : X \rightarrow Y$ in \mathcal{M} , we have $\gamma f_0 = \gamma f_1$ if and only if f_0 and f_1 are homotopic.*

Proof. As noted in [remark 3.2.4](#), if $f_0, f_1 : X \rightarrow Y$ are homotopic, then we must have $\gamma f_0 = \gamma f_1$. Conversely, suppose $\gamma f_0 = \gamma f_1$ with X cofibrant and Y fibrant. Let (RX, i') be a cofibrant fibrant replacement for X and (QY, p') be a fibrant cofibrant replacement for Y . Then, there exists morphisms $f'_0, f'_1 : RX \rightarrow QY$ such that $f_0 = p' \circ f'_0 \circ i'$ and $f_1 = p' \circ f'_1 \circ i'$. Since $i' : X \rightarrow RX$ and $p' : QY \rightarrow Y$ are weak equivalences, we must have $\gamma f'_0 = \gamma f'_1$ in $\text{Ho } \mathcal{M}$. The theorem then implies f'_0 and f'_1 are homotopic; thus f_0 and f_1 are also homotopic, by lemmas [3.2.8](#) and [3.2.10](#). ■

GENERALITIES

A.I Factorisation systems

Definition A.I.1. Let $f : X \rightarrow Y$ and $g : Z \rightarrow W$ be morphisms in a category \mathcal{C} . Given a commutative square in \mathcal{C} ,

$$\begin{array}{ccc} Z & \xrightarrow{z} & X \\ g \downarrow & & \downarrow f \\ W & \xrightarrow{w} & Y \end{array}$$

a **lift** is a morphism $h : W \rightarrow X$ such that $f \circ h = w$ and $h \circ g = z$.

We say g has the **left lifting property** with respect to f and f has the **right lifting property** with respect to g if every commutative square in \mathcal{C} of the form above has a lift; and we say f is **left orthogonal** to g and g is **right orthogonal** to f if lifts exist *and* are unique.

Lemma A.I.2. Let $f : X \rightarrow Y$ and $g : Z \rightarrow W$ be morphisms in a locally small category \mathcal{C} . Consider the commutative diagram in **Set** shown below,

$$\begin{array}{ccccc} \mathcal{C}(W, X) & & & & \mathcal{C}(W, Y) \\ & \searrow^{f_*} & & & \downarrow g_* \\ & & \bullet & \longrightarrow & \mathcal{C}(W, Y) \\ & \searrow^{g_*} & \downarrow & & \downarrow g_* \\ & & \mathcal{C}(Z, X) & \xrightarrow{f_*} & \mathcal{C}(Z, Y) \end{array}$$

where the inner square is a pullback diagram.

- (i) The dashed arrow is a surjection if and only if g has the left lifting property with respect to f .
- (ii) The dashed arrow is a bijection if and only if g is left orthogonal to f .

Proof. This is just a restatement of the definition. ■

Lemma A.1.3. *Let $f : X \rightarrow Y$ be a morphism in a category \mathcal{C} . The following are equivalent:*

- (i) f is an isomorphism.
- (ii) f is right orthogonal to any morphism in \mathcal{C} .
- (iii) f has the right lifting property with respect to any morphism in \mathcal{C} .
- (iv) f has the right lifting property with respect to itself.

Dually, the following are equivalent:

- (i') f is an isomorphism.
- (ii') f is left orthogonal to any morphism in \mathcal{C} .
- (iii') f has the left lifting property with respect to any morphism in \mathcal{C} .
- (iv') f has the left lifting property with respect to itself.

Proof. (i) \Rightarrow (ii). Suppose $r : Y \rightarrow X$ is a morphism such that $r \circ f = \text{id}_X$. Then, for any commutative square as below,

$$\begin{array}{ccc} Z & \xrightarrow{z} & X \\ g \downarrow & & \downarrow f \\ W & \xrightarrow{w} & Y \end{array}$$

we have $(r \circ w) \circ g = r \circ f \circ z = z$; but if $f \circ r = \text{id}_Y$ as well, then $f \circ (r \circ w) = w$; thus $r \circ w : W \rightarrow X$ is the required lift. It is clearly unique, as f is monic.

(ii) \Rightarrow (iii), (iii) \Rightarrow (iv). Obvious.

(iv) \Rightarrow (i). Consider the following commutative square:

$$\begin{array}{ccc} X & \xrightarrow{\text{id}} & X \\ f \downarrow & & \downarrow f \\ Y & \xrightarrow{\text{id}} & Y \end{array}$$

Since f has the right lifting property with respect to itself, there exists a morphism $h : Y \rightarrow X$ such that $h \circ f = \text{id}_X$ and $f \circ h = \text{id}_Y$. ■

Definition A.I.4. A **weak factorisation system** for a category \mathcal{C} is a pair $(\mathcal{L}, \mathcal{R})$ of subclasses of $\text{mor } \mathcal{C}$ satisfying these conditions:

- For each morphism f in \mathcal{C} there exists a pair (g, h) with $g \in \mathcal{L}$ and $h \in \mathcal{R}$ such that $f = h \circ g$. Such a pair is a **$(\mathcal{L}, \mathcal{R})$ -factorisation** of f .
- A morphism is in \mathcal{L} if and only if it has the left lifting property with respect to every morphism in \mathcal{R} .
- A morphism is in \mathcal{R} if and only if it has the right lifting property with respect to every morphism in \mathcal{L} .

An **orthogonal factorisation system** is the same thing as a weak factorisation system, except for replacing ‘... has the left/right lifting property with respect to ...’ with ‘... is left/right orthogonal to ...’.

Remark A.I.5. Obviously, $(\mathcal{L}, \mathcal{R})$ is a weak (resp. orthogonal) factorisation system for \mathcal{C} if and only if $(\mathcal{R}^{\text{op}}, \mathcal{L}^{\text{op}})$ is a weak (resp. orthogonal) factorisation system for \mathcal{C}^{op} .

Lemma A.I.6. *Let A be an object in a category \mathcal{C} with a weak (resp. orthogonal) factorisation system $(\mathcal{L}, \mathcal{R})$. Then the slice category $\mathcal{C}_{/A}$ has a weak (resp. orthogonal) factorisation system where a morphism is in the left or right class if and only if it is so in \mathcal{C} .*

Proof. The projection $\mathcal{C}_{/A} \rightarrow \mathcal{C}$ induces a bijection between solutions for lifting problems in $\mathcal{C}_{/A}$ and solutions for the corresponding lifting problems in \mathcal{C} . ■

Proposition A.I.7. *Let $(\mathcal{L}, \mathcal{R})$ be a weak or orthogonal factorisation system for a category \mathcal{C} .*

(i) Given a pullback diagram in \mathcal{C} as below,

$$\begin{array}{ccc} X' & \xrightarrow{p} & X \\ f' \downarrow & & \downarrow f \\ Y' & \xrightarrow{q} & Y \end{array}$$

if the morphism f is in \mathcal{R} , then f' is also in \mathcal{R} .

(ii) Let I be a set. If $f_i : X_i \rightarrow Y_i$ is a morphism in \mathcal{R} for all i in I and the product $\prod_i f_i : \prod_i X_i \rightarrow \prod_i Y_i$ exists in \mathcal{C} , then $\prod_i f_i$ is also in \mathcal{R} .

(iii) Given a commutative diagram of the form

$$\begin{array}{ccccc} X' & \xrightarrow{i_X} & X & \xrightarrow{r_X} & X' \\ f' \downarrow & & f \downarrow & & \downarrow f' \\ Y' & \xrightarrow{i_Y} & Y & \xrightarrow{r_Y} & Y' \end{array}$$

where $r_X \circ i_X = \text{id}_{X'}$ and $r_Y \circ i_Y = \text{id}_{Y'}$, if f is in \mathcal{R} , then so is f' ; in other words, \mathcal{R} is closed under retracts.

(iv) \mathcal{L} is closed under composition.

(v) Let γ be an ordinal and let $Z : \gamma \rightarrow \mathcal{C}$ be a functor that preserves sequential colimits. We write Z_α for $Z(\alpha)$, where $\alpha < \gamma$, and $g_{\alpha,\beta} : Z_\alpha \rightarrow Z_\beta$ for the morphism $Z(\alpha \rightarrow \beta)$, where $\alpha < \beta < \gamma$. If λ is a colimiting cocone from Z to W and each $g_{\alpha,\beta}$ is in \mathcal{L} , then each component $\lambda_\alpha : Z_\alpha \rightarrow W$ is also in \mathcal{L} .

Proof. (i). Suppose g is in \mathcal{L} and consider the following commutative diagram:

$$\begin{array}{ccccc} Z & \xrightarrow{z} & X' & \xrightarrow{p} & X \\ g \downarrow & & \downarrow f' & & \downarrow f \\ W & \xrightarrow{w} & Y' & \xrightarrow{q} & Y \end{array}$$

There exists $h : Z \rightarrow X$ such that $h \circ g = p \circ z$ and $f \circ h = q \circ w$. In particular, there exists a unique morphism $h' : Z \rightarrow X'$ such that $f' \circ h' = w$ and $p \circ h' = h$, by the universal property of pullbacks. Thus $p \circ h' \circ g = h \circ g = p \circ z$ and

$f' \circ h' \circ g = w \circ g = f' \circ z$, but p and f' are jointly monic, so $h' \circ g = z$. Thus we have the required lift, and h' is unique if h is.

(ii). We may construct the required lift componentwise.

(iii). Suppose g is in \mathcal{L} and consider the following commutative diagram:

$$\begin{array}{ccccccc}
 Z & \xrightarrow{z} & X' & \xrightarrow{i_X} & X & \xrightarrow{r_X} & X' \\
 g \downarrow & & \downarrow f' & & \downarrow f & & \downarrow f' \\
 W & \xrightarrow{w} & Y' & \xrightarrow{i_Y} & Y & \xrightarrow{r_Y} & Y'
 \end{array}$$

There exists $h : Z \rightarrow X$ such that $h \circ g = i_X \circ z$ and $f \circ h = i_Y \circ w$, and so for $h' = r_X \circ h$:

$$\begin{aligned}
 h' \circ g &= r_X \circ i_X \circ z = z \\
 f' \circ h' &= f' \circ r_X \circ h = r_Y \circ f \circ h = r_Y \circ i_Y \circ w = w
 \end{aligned}$$

Thus $h' : Z \rightarrow X'$ is the required lift, and h' is unique if h is (because i_X is split monic).

(iv). Suppose $g' : Z' \rightarrow Z$ and $g : Z \rightarrow W$ are in \mathcal{L} and $f : X \rightarrow Y$ is in \mathcal{R} . Consider the following commutative diagram:

$$\begin{array}{ccc}
 Z' & \xrightarrow{z'} & X \\
 g' \downarrow & & \downarrow f \\
 Z & & \\
 g \downarrow & & \downarrow \\
 W & \xrightarrow{w} & Y
 \end{array}$$

There must exist a morphism $z : Z \rightarrow X$ such that $z \circ g' = z'$ and $f \circ z' = w \circ g$, and hence a morphism $h : W \rightarrow X$ such that $h \circ g = z$ and $f \circ h = w$. Obviously, $h \circ (g' \circ g) = z'$, so h is the required lift and is moreover unique if $(\mathcal{L}, \mathcal{R})$ is an orthogonal factorisation system.

(v). We may assume without loss of generality that $\alpha = 0$, since any non-empty terminal segment of γ is cofinal in γ . Suppose $f : X \rightarrow Y$ is in \mathcal{R} and consider

the following commutative diagram:

$$\begin{array}{ccc} Z_0 & \xrightarrow{z_0} & X \\ \lambda_0 \downarrow & & \downarrow f \\ W & \xrightarrow{w} & Y \end{array}$$

For each $\alpha < \gamma$, given z_α making the following diagram commute,

$$\begin{array}{ccccc} Z_\alpha & \xrightarrow{z_\alpha} & & \longrightarrow & X \\ g_{\alpha,\alpha+1} \downarrow & & & & \downarrow f \\ Z_{\alpha+1} & \xrightarrow{\lambda_{\alpha+1}} & W & \xrightarrow{w} & Y \end{array}$$

choose a lift $z_{\alpha+1} : Z_{\alpha+1} \rightarrow X$; for each limit ordinal $\beta < \gamma$, let $z_\beta : Z_\beta \rightarrow X$ be the unique morphism such that $z_\beta \circ g_{\alpha,\beta} = z_\alpha$ for all $\alpha < \beta$. (Such z_β exist and are unique because $Z_\beta = \varinjlim_{\alpha < \beta} Z_\alpha$.) Note that the universal property of W then guarantees that $w \circ \lambda_\beta = f \circ z_\beta$.

Having constructed morphisms $z_\alpha : Z_\alpha \rightarrow X$ for all $\alpha < \gamma$ as above, we may now obtain $h : W \rightarrow X$ as the unique morphism such that $h \circ \lambda_\alpha = z_\alpha$ for all $\alpha < \gamma$, and again we automatically have $f \circ h = w$. It is also clear that h is unique if $(\mathcal{L}, \mathcal{R})$ is an orthogonal factorisation system. \blacksquare

Proposition A.1.8. *Let $(\mathcal{L}, \mathcal{R})$ be a weak (resp. orthogonal) factorisation system for a category \mathcal{C} , and let $(\mathcal{L}', \mathcal{R}')$ be a weak (resp. orthogonal) factorisation system for a category \mathcal{C}' . Given an adjunction*

$$F \dashv U : \mathcal{C}' \rightarrow \mathcal{C}$$

the following are equivalent:

- (i) *F sends morphisms in \mathcal{L} to morphisms in \mathcal{L}' .*
- (ii) *U sends morphisms in \mathcal{R}' to morphisms in \mathcal{R} .*

Proof. The adjunction induces a bijection between solutions to the two lifting problems shown below:

$$\begin{array}{ccc} FZ & \longrightarrow & X \\ Fg \downarrow & \nearrow ? & \downarrow f \\ FW & \longrightarrow & Y \end{array} \qquad \begin{array}{ccc} Z & \longrightarrow & UX \\ g \downarrow & \nearrow ? & \downarrow Uf \\ W & \longrightarrow & UY \end{array}$$

Thus, Fg has the left lifting property (resp. is left orthogonal) with respect to f if and only if Uf has the right lifting property (resp. is right orthogonal) with respect to g . ■

Definition A.I.9. A **functorial factorisation system** on a category \mathcal{C} is a pair of functors $L, R : [\mathbb{2}, \mathcal{C}] \rightarrow [\mathbb{2}, \mathcal{C}]$ satisfying the following equations:

$$\text{dom} \circ L = \text{dom} \quad \text{codom} \circ L = \text{dom} \circ R \quad \text{codom} \circ R = \text{codom}$$

Here, dom and codom are considered as functors $[\mathbb{2}, \mathcal{C}] \rightarrow \mathcal{C}$. A **functorial weak** (resp. **orthogonal**) **factorisation system** on \mathcal{C} is a weak (resp. orthogonal) factorisation system $(\mathcal{L}, \mathcal{R})$ together with a functorial factorisation system (L, R) such that $Lf \in \mathcal{L}$ and $Rf \in \mathcal{R}$ for all morphisms f in \mathcal{C} .

Proposition A.I.10. Any orthogonal factorisation system can be extended to a functorial one.

Proof. For each morphism f in a category \mathcal{C} with an orthogonal factorisation system $(\mathcal{L}, \mathcal{R})$, choose a factorisation $f = Rf \circ Lf$ with $Lf \in \mathcal{L}$ and $Rf \in \mathcal{R}$. Given a commutative square in \mathcal{C} , say

$$\begin{array}{ccc} Z & \xrightarrow{g} & W \\ z \downarrow & & \downarrow w \\ X & \xrightarrow{f} & Y \end{array}$$

the lifting property ensures that the dashed arrow in the diagram below exists,

$$\begin{array}{ccccc} Z & \xrightarrow{Lg} & \bullet & \xrightarrow{Rg} & W \\ z \downarrow & & \vdots & & \downarrow w \\ X & \xrightarrow{Lf} & \bullet & \xrightarrow{Rf} & Y \end{array}$$

and orthogonality ensures uniqueness and hence functoriality. ■

Proposition A.I.11. Let A be an object in a category \mathcal{C} and let $\Sigma_A : \mathcal{C}_{/A} \rightarrow \mathcal{C}$ be the projection from the slice category.

- (i) For each functorial factorisation system (L, R) on \mathcal{C} , there exists a unique functorial factorisation system (L_A, R_A) on $\mathcal{C}_{/A}$ such that

$$[\mathbb{2}, \Sigma_A] \circ L_A = L \circ [\mathbb{2}, \Sigma_A] \quad [\mathbb{2}, \Sigma_A] \circ R_A = R \circ [\mathbb{2}, \Sigma_A]$$

where $[\mathbb{2}, \Sigma_A] : [\mathbb{2}, \mathcal{C}_{/A}] \rightarrow [\mathbb{2}, \mathcal{C}]$ is the evident induced functor.

- (ii) If (L, R) is part of a functorial weak or orthogonal factorisation system on C , then (L_A, R_A) is compatible with the induced weak or orthogonal factorisation system on $C_{/A}$ as well.

Proof. Obvious. ■

A.2 Relative categories

Prerequisites. § 0.1.

In this section we use the explicit universe convention.

Definition A.2.1. A **relative category** C consists of a category $\text{und } C$ and a subcategory $\text{weq } C$ such that $\text{ob } \text{und } C = \text{ob } \text{weq } C$. We say $\text{und } C$ is the **underlying category** of C , and that the morphisms in $\text{weq } C$ are the **weak equivalences** in C .

Remark A.2.2. The subcategory $\text{weq } C$ is entirely determined by $\text{mor } \text{weq } C$, so a relative category may equivalently be defined as a category equipped with a distinguished subset of morphisms closed under composition and containing all the identity morphisms.

For brevity, we will write $\text{ob } C$ for $\text{ob } \text{und } C$, $\text{mor } C$ for $\text{mor } \text{und } C$, and we may occasionally abuse notation and write $\text{weq } C$ instead of $\text{mor } \text{weq } C$.

Remark A.2.3. Every category C can be endowed with the structure of a relative category in two ways: we can make it into a **minimal relative category** $\text{min } C$ by taking $\text{weq } \text{min } C$ to be the set of identity morphisms in C ; or we could make it into a **maximal relative category** $\text{max } C$ by taking $\text{weq } \text{max } C = \text{mor } C$. We may also define the **minimal saturated relative category** $\text{min}^+ C$ by taking $\text{weq } \text{min}^+ C$ to be the set of all isomorphisms in C .

Definition A.2.4. Given a relative category C , the **opposite relative category** C^{op} is defined by $\text{und } C^{\text{op}} = (\text{und } C)^{\text{op}}$ and $\text{weq } C^{\text{op}} = (\text{weq } C)^{\text{op}}$.

Definition A.2.5. Let C and D be relative categories. A **relative functor** $C \rightarrow D$ is a functor $\text{und } C \rightarrow \text{und } D$ that sends weak equivalences in C to weak equivalences in D . The **relative functor category** $[C, D]_{\text{h}}$ is the full subcategory of $[\text{und } C, \text{und } D]$ spanned by the relative functors, and the weak equivalences in $[C, D]_{\text{h}}$ are defined to be the natural transformations that are componentwise weak equivalences in D .

Definition A.2.6. Let C be a category and let $\mathcal{W} \subseteq \text{mor } C$. A **localisation of C away from \mathcal{W}** is a category $C[\mathcal{W}^{-1}]$ equipped with a functor $\gamma : C \rightarrow C[\mathcal{W}^{-1}]$ with the following universal property:

- Given a functor $F : C \rightarrow D$ such that Ff is an isomorphism for all f in \mathcal{W} , there exists a unique functor $\overline{F} : C[\mathcal{W}^{-1}] \rightarrow D$ such that $\overline{F}\gamma = F$.

Remark A.2.7. The universal property in the above definition is strict; as such, $C[\mathcal{W}^{-1}]$ is unique up to unique isomorphism. Nonetheless, $C[\mathcal{W}^{-1}]$ automatically has a 2-universal property: if $F, G : C \rightarrow D$ both factor through $C[\mathcal{W}^{-1}]$, then so do all natural transformations $F \Rightarrow G$.

Proposition A.2.8. *If C is a \mathbf{U} -small category, then there exists a \mathbf{U} -small category with the universal property of $C[\mathcal{W}^{-1}]$.*

Proof. Use the general adjoint functor theorem. ■

Definition A.2.9. The **homotopy category** of a relative category C is a localisation of $\text{und } C$ away from $\text{weq } C$ and is denoted $\text{Ho } C$. A **semi-saturated relative category** is a relative category in which every isomorphism is a weak equivalence. A **saturated relative category** is a relative category C such that the weak equivalences in C are precisely the ones that become isomorphisms in $\text{Ho } C$.

Remark A.2.10. Obviously, there is no loss of generality in considering semi-saturated relative categories and their homotopy categories instead of localisations $C[\mathcal{W}^{-1}]$ for arbitrary subsets $\mathcal{W} \subseteq \text{mor } C$.

Remark A.2.11. Clearly, every saturated relative category is semi-saturated, and a minimal saturated relative category is indeed saturated in the sense above.

Definition A.2.12. Let C be a category and let \mathcal{W} be a subset of $\text{mor } C$. The **2-out-of-3 property** for \mathcal{W} says:

- Given any two morphisms $f : X \rightarrow Y, g : Y \rightarrow Z$ in C , if any two of f, g , or $g \circ f$ are in \mathcal{W} , then all of them are.

The **2-out-of-6 property** for \mathcal{W} says:

- Given any three morphisms $f : X \rightarrow Y, g : Y \rightarrow Z, h : Y \rightarrow Z$ in C , if both $h \circ g$ and $g \circ f$ are in \mathcal{W} , then so too are f, g, h , and $h \circ g \circ f$.

Lemma A.2.13. *Let C be a category and let $\mathcal{W} \subseteq \text{mor } C$.*

- (i) If \mathcal{W} has the 2-out-of-6 property, then it also has the 2-out-of-3 property.
- (ii) The set of all isomorphisms in C has the 2-out-of-6 property.
- (iii) If $F : C' \rightarrow C$ is a functor and \mathcal{W} has either the 2-out-of-3 property or the 2-out-of-6 property, then $F^{-1}\mathcal{W}$ has the same property.

Proof. (i). Consider the three cases $f = \text{id}$, $g = \text{id}$, $h = \text{id}$ in turn.

(ii). If $h \circ g$ and $g \circ f$ are isomorphisms, then g must be split epic and split monic; thus g itself is an isomorphism, hence so too are f and h .

(iii). Obvious. ■

Corollary A.2.14. *If C is a saturated relative category, then $\text{weq } C$ has the 2-out-of-6 property.* ■

Lemma A.2.15. *Let \mathbf{RelCat} be the category of \mathbf{U} -small relative categories and relative functors, let $\mathbf{SsRelCat}$ be the full subcategory of semi-saturated relative categories, and let \mathbf{Cat} be the category of \mathbf{U} -small categories and functors.*

- (i) \mathbf{RelCat} has finite products and is cartesian closed, where the product of C and D is the cartesian product $C \times D$ with weak equivalences taken componentwise, and the exponential of \mathcal{E} by D is the relative functor category $[D, \mathcal{E}]_h$; moreover, $\mathbf{SsRelCat}$ is closed under finite products in \mathbf{RelCat} , and if \mathcal{E} is semi-saturated, then so is $[D, \mathcal{E}]_h$.
- (ii) If D is an ordinary category and \mathcal{E} is a relative category, then the inclusion

$$\text{und } [\min D, \mathcal{E}]_h \hookrightarrow [D, \text{und } \mathcal{E}]$$

is an isomorphism of ordinary categories, natural in D and \mathcal{E} ; and if \mathcal{E} is moreover semi-saturated, then the inclusion

$$\text{und } [\min^+ D, \mathcal{E}]_h \hookrightarrow [D, \text{und } \mathcal{E}]$$

is also an isomorphism of ordinary categories, natural in D and \mathcal{E} .

- (iii) If D is a relative category and \mathcal{E} is an ordinary category, then the canonical relative functor

$$\min^+ [\text{Ho } D, \mathcal{E}] \rightarrow [D, \min^+ \mathcal{E}]_h$$

is an isomorphism of relative categories, natural in \mathcal{D} and \mathcal{E} ; in particular, the \mathbf{U} -small minimal saturated relative categories form an exponential ideal in \mathbf{RelCat} .

- (iv) The functors $\min : \mathbf{Cat} \rightarrow \mathbf{RelCat}$ and $\min^+ : \mathbf{Cat} \rightarrow \mathbf{SsRelCat}$ are cartesian closed functors, i.e. they preserve finite products, and the canonical relative functors

$$\begin{aligned} \min [\mathcal{D}, \mathcal{E}] &\rightarrow [\min \mathcal{D}, \min \mathcal{E}]_{\mathbf{h}} \\ \min^+ [\mathcal{D}, \mathcal{E}] &\rightarrow [\min^+ \mathcal{D}, \min^+ \mathcal{E}]_{\mathbf{h}} \end{aligned}$$

are isomorphisms of relative categories, natural in \mathcal{D} and \mathcal{E} .

- (v) If \mathcal{C} and \mathcal{D} are relative categories, then the canonical functor

$$\mathrm{Ho}(\mathcal{C} \times \mathcal{D}) \rightarrow \mathrm{Ho}(\mathcal{C}) \times \mathrm{Ho}(\mathcal{D})$$

is an isomorphism of ordinary categories, natural in \mathcal{C} and \mathcal{D} .

Proof. Claims (i) – (iv) are straightforward from the definitions. For claim (v), we simply observe that we have a chain of isomorphisms

$$\begin{aligned} \mathrm{Fun}(\mathrm{Ho}(\mathcal{C}) \times \mathrm{Ho}(\mathcal{D}), \mathcal{E}) &\cong \mathrm{Fun}(\mathrm{Ho} \mathcal{C}, [\mathrm{Ho} \mathcal{D}, \mathcal{E}]) \\ &\cong \mathrm{RelFun}(\mathcal{C}, \min^+ [\mathrm{Ho} \mathcal{D}, \mathcal{E}]) \\ &\cong \mathrm{RelFun}(\mathcal{C}, [\mathcal{D}, \min^+ \mathcal{E}]_{\mathbf{h}}) \\ &\cong \mathrm{RelFun}(\mathcal{C} \times \mathcal{D}, \min^+ \mathcal{E}) \\ &\cong \mathrm{Fun}(\mathrm{Ho}(\mathcal{C} \times \mathcal{D}), \mathcal{E}) \end{aligned}$$

natural in \mathcal{C} , \mathcal{D} , and \mathcal{E} , so by taking $\mathcal{E} = \mathrm{Ho}(\mathcal{C}) \times \mathrm{Ho}(\mathcal{D})$ and $\mathcal{E} = \mathrm{Ho}(\mathcal{C} \times \mathcal{D})$ in turn, we see that Ho preserves binary products. \blacksquare

Proposition A.2.16. *Let $\mathbf{SsRelCat}$ be the category of \mathbf{U} -small semi-saturated relative categories and relative functors, and let \mathbf{Cat} be the category of \mathbf{U} -small categories and functors. There is then a string of adjoint functors*

$$\mathrm{Ho} \dashv \min^+ \dashv \mathrm{und} \dashv \max \dashv \mathrm{weq} : \mathbf{SsRelCat} \rightarrow \mathbf{Cat}$$

where Ho sends a relative category to its homotopy category, \min^+ makes an ordinary category into a minimal saturated relative category, und sends a relative category to its underlying category, \max makes an ordinary category into a

maximal relative category, and weq sends a relative category to its subcategory of weak equivalences. Moreover, both min^+ and max are fully faithful, and Ho preserves finite products.


Proof. Obvious. ■

Definition A.2.17. A **zigzag type** is a relative category T where $\text{und } T$ is the free category on an inhabited finite planar graph of the form



where the edges are arrows that point either left or right, and $\text{weq } T$ consists of all identities and all composites of left-pointing arrows. A **morphism of zigzag types** is a relative functor that maps the leftmost object to the leftmost object and the rightmost object to the rightmost object. We write \mathbf{T} for the category of zigzag types.^[1]

A **zigzag** of type T in a relative category C is a relative functor $T \rightarrow C$. Given objects X and Y in C , we denote by $C^T(X, Y)$ the category whose objects are the zigzags starting at X and ending at Y and whose morphisms are commutative diagrams in C of the form



where the rows are zigzags of type T and the unmarked columns are weak equivalences.

Remark A.2.18. It is clear that $C^T(X, Y)$ is a subcategory of the relative functor category $[T, C]_{\text{h}}$. Thus, if C is a \mathbf{U} -small relative category, precomposition makes the assignment $T \mapsto C^T(X, Y)$ into a functor $\mathbf{T}^{\text{op}} \rightarrow \mathbf{Cat}$, which we denote by $C^*(X, Y)$. The Grothendieck construction applied to this functor yields the following \mathbf{U} -small category $C^{(\mathbf{T})}(X, Y)$:

- Its objects are pairs (T, f) , where T is a zigzag type and f is a zigzag of type T in C .

^[1] Warning: This is the *opposite* of the category \mathbf{T} defined in [DHKS, § 34].

- A morphism $(T', f') \rightarrow (T, f)$ is a pair (α, β) where $\alpha : T' \rightarrow T$ is a morphism in \mathbf{T} and $\beta : f' \rightarrow \alpha^* f$ is a morphism in $\mathcal{C}^{T'}(X, Y)$.
- The composite of a pair of morphisms $(\alpha', \beta') : (T'', f'') \rightarrow (T', f')$ and $(\alpha, \beta) : (T', f') \rightarrow (T, f)$ is given by $(\alpha \circ \alpha', \alpha'^* \beta \circ \beta')$.

There is an evident projection functor $\mathcal{C}^{(\mathbf{T})}(X, Y) \rightarrow \mathbf{T}$, and by construction it is a Grothendieck fibration with a canonical splitting.

Example A.2.19. If $f : X \rightarrow Y$ is a weak equivalence in a relative category \mathcal{C} , then we have commutative diagrams

$$\begin{array}{ccc}
X & \xrightarrow{\text{id}} & X & \xleftarrow{\text{id}} & X \\
\text{id} \downarrow & & f \downarrow & & \downarrow \text{id} \\
X & \xrightarrow{f} & Y & \xleftarrow{f} & X
\end{array}
\qquad
\begin{array}{ccccc}
Y & \xleftarrow{f} & X & \xrightarrow{f} & Y \\
\text{id} \downarrow & & f \downarrow & & \downarrow \text{id} \\
Y & \xleftarrow{\text{id}} & Y & \xrightarrow{\text{id}} & Y
\end{array}$$

and these correspond to morphisms of zigzags in \mathcal{C} .

Theorem A.2.20. Let X and Y be objects in a relative category \mathcal{C} .

- For each zigzag type T , the map that sends an object in $\mathcal{C}^T(X, Y)$ to the corresponding composite in $\text{Ho } \mathcal{C}(X, Y)$ is a functor when the latter is regarded as a discrete category.
- The functors described above constitute a jointly surjective cocone from the diagram $\mathcal{C}^*(X, Y)$ to $\text{Ho } \mathcal{C}(X, Y)$.
- The induced functor $\mathcal{C}^{(\mathbf{T})}(X, Y) \rightarrow \text{Ho } \mathcal{C}(X, Y)$ is surjective, and moreover two objects in $\mathcal{C}^{(\mathbf{T})}(X, Y)$ become equal in $\text{Ho } \mathcal{C}$ if and only if they are in the same connected component.

Proof. All obvious except for the last part of claim (iii), for which we refer to paragraphs 33.8 and 33.10 in [DHKS]. ■

Definition A.2.21. Two objects in a relative category are **weakly equivalent** if they can be connected by a *zigzag* of weak equivalences.

Remark A.2.22. If X and Y are weakly equivalent in a relative category \mathcal{C} , then they are isomorphic in $\text{Ho } \mathcal{C}$.

A.3 Kan extensions

Prerequisites. § 0.1.

In this section we use the explicit universe convention.

Definition A.3.1. Let $F : C \rightarrow D$ and $G : C \rightarrow \mathcal{E}$ be two functors. A **left Kan extension** (resp. **right Kan extension**) of G along F is an initial (resp. terminal) object of the category $(G \downarrow F^*)$ (resp. $(F^* \downarrow G)$) described below:

- The objects are pairs (H, α) where H is a functor $D \rightarrow \mathcal{E}$ and α is a natural transformation of type $G \Rightarrow HF$ (resp. $HF \Rightarrow G$).
- The morphisms $(H', \alpha') \rightarrow (H, \alpha)$ are those natural transformations $\beta : H' \Rightarrow H$ such that $\beta F \bullet \alpha' = \alpha$ (resp. $\alpha \bullet \beta F = \alpha'$).

¶ **A.3.2.** Clearly, Kan extensions are unique up to unique isomorphism if they exist. We write $(\text{Lan}_G F, \eta)$ for the left Kan extension of F along G , and $(\text{Ran}_G F, \varepsilon)$ for the right Kan extension of F along G .

Definition A.3.3. Let $F : C \rightarrow D$, $G : C \rightarrow \mathcal{E}$, and $L : \mathcal{E} \rightarrow \mathcal{F}$ be three functors. We say L **preserves** a left (resp. right) Kan extension (H, α) of F along G if $(KH, K\alpha)$ is a left (resp. right) Kan extension of KF along G .

Let **Set** be the category of **U**-small sets, and suppose \mathcal{E} is locally **U**-small. We say a left Kan extension $(\text{Lan}_G F, \eta)$ is **pointwise** if it is preserved by all functors of the form $\mathcal{E}(-, E) : \mathcal{E} \rightarrow \mathbf{Set}^{\text{op}}$; dually, we say a right Kan extension $(\text{Ran}_G F, \varepsilon)$ is **pointwise** if it is preserved by all functors of the form $\mathcal{E}(E, -) : \mathcal{E} \rightarrow \mathbf{Set}$. If a Kan extension is preserved by *all* functors, then it is said to be **absolute**.

BIBLIOGRAPHY

Adámek, J. and J. Rosický

[LPAC] *Locally presentable and accessible categories*. London Mathematical Society Lecture Note Series 189. Cambridge: Cambridge University Press, 1994. xiv+316. ISBN: 0-521-42261-2. DOI: [10.1017/CB09780511600579](https://doi.org/10.1017/CB09780511600579).

Borceux, Francis

[1994a] *Handbook of categorical algebra. 1. Basic category theory*. Encyclopedia of Mathematics and its Applications 50. Cambridge: Cambridge University Press, 1994. xvi+345. ISBN: 0-521-44178-1.

[1994b] *Handbook of categorical algebra. 2. Categories and structures*. Encyclopedia of Mathematics and its Applications 51. Cambridge: Cambridge University Press, 1994. xviii+443. ISBN: 0-521-44179-X.

Dugger, Daniel

[2001] ‘Universal homotopy theories’. In: *Adv. Math.* 164.1 (2001), pp. 144–176. ISSN: 0001-8708. DOI: [10.1006/aima.2001.2014](https://doi.org/10.1006/aima.2001.2014).

Dwyer, William G., Philip S. Hirschhorn, Daniel M. Kan and Jeffrey H. Smith

[DHKS] *Homotopy limit functors on model categories and homotopical categories*. Vol. 113. Mathematical Surveys and Monographs. Providence, RI: American Mathematical Society, 2004. viii+181. ISBN: 0-8218-3703-6.

Dwyer, William G. and Jan Spaliski

[DS] ‘Homotopy theories and model categories’. In: *Handbook of algebraic topology*. Amsterdam: North-Holland, 1995, pp. 73–126. DOI: [10.1016/B978-044481779-2/50003-1](https://doi.org/10.1016/B978-044481779-2/50003-1).

BIBLIOGRAPHY

Gabriel, Peter and Friedrich Ulmer

[1971] *Lokal präsentierbare Kategorien*. Lecture Notes in Mathematics 221. Berlin: Springer-Verlag, 1971. v+200.

Gabriel, Peter and Michel Zisman

[GZ] *Calculus of fractions and homotopy theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35. Springer-Verlag New York, Inc., New York, 1967. x+168.

Goerss, Paul G. and John F. Jardine

[GJ] *Simplicial homotopy theory*. Vol. 174. Progress in Mathematics. Basel: Birkhäuser Verlag, 1999. xvi+510. ISBN: 3-7643-6064-X. DOI: [10.1007/978-3-0348-8707-6](https://doi.org/10.1007/978-3-0348-8707-6).

Hirschhorn, Philip S.

[2003] *Model categories and their localizations*. Vol. 99. Mathematical Surveys and Monographs. Providence, RI: American Mathematical Society, 2003. xvi+457. ISBN: 0-8218-3279-4.

Hovey, Mark

[1999] *Model categories*. Vol. 63. Mathematical Surveys and Monographs. Providence, RI: American Mathematical Society, 1999. xii+209. ISBN: 0-8218-1359-5.

Mac Lane, Saunders

[CWM] *Categories for the working mathematician*. Second. Graduate Texts in Mathematics 5. New York: Springer-Verlag, 1998. xii+314. ISBN: 0-387-98403-8.

Mac Lane, Saunders and Ieke Moerdijk

[ML–M] *Sheaves in geometry and logic. A first introduction to topos theory*. Corrected. Universitext. New York: Springer-Verlag, 1992. xii+629. ISBN: 0-387-97710-4.

Mathias, Adrian R. D.

[2001] ‘The strength of Mac Lane set theory’. In: *Ann. Pure Appl. Logic* 110.1-3 (2001), pp. 107–234. ISSN: 0168-0072. DOI: [10.1016/S0168-0072\(00\)00031-2](https://doi.org/10.1016/S0168-0072(00)00031-2).

May, J. Peter

[1967] *Simplicial objects in algebraic topology*. Van Nostrand Mathematical Studies, No. 11. Princeton, NJ: D. Van Nostrand Co., Inc., 1967. vi+161.

Munkres, James R.

[2000] *Topology*. English. Second. Upper Saddle River, NJ: Prentice Hall, 2000. xvi+537. ISBN: 0-13-181629-2.

Quillen, Daniel G.

[1967] *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Berlin: Springer-Verlag, 1967. iv+156.

[1969] 'Rational homotopy theory'. In: *Ann. of Math. (2)* 90 (1969), pp. 205–295. ISSN: 0003-486X.

Shulman, Michael A.

[2012] 'Exact completions and small sheaves'. In: *Theory Appl. Categ.* 27 (2012), pp. 97–173. ISSN: 1201-561X.

Waterhouse, William C.

[1975] 'Basically bounded functors and flat sheaves'. In: *Pacific J. Math.* 57.2 (1975), pp. 597–610. ISSN: 0030-8730.

Weibel, Charles A.

[1994] *An introduction to homological algebra*. Cambridge Studies in Advanced Mathematics 38. Cambridge: Cambridge University Press, 1994. xiv+450. ISBN: 0-521-43500-5.

INDEX

- 2-out-of-3 property, 65
- 2-out-of-6 property, 65
- accessible category, 13
 - classification theorem, 15
- accessible extension, 26, 28, 30
- accessible functor, 15
- adjoint functor theorem
 - accessible —, 21, 29
- \aleph -number, 7
- anodyne extension, 40
- arity class, 8
 - classification theorem, 8
- Beck–Chevalley condition, 24
- cardinal, 7
 - classification theorem, 7
 - regular —, 8
 - strongly inaccessible —, 9
- cartesian closed category, 38
- category
 - finite —, 8
 - locally small —, 2
 - small —, 2, 8
- category with weak equivalences, 41
- class, 2
- classification theorem
 - for accessible categories, 15
 - for arity classes, 8
 - for cardinalities, 7
 - for locally presentable categories, 17
 - for well-ordered sets, 5
- cocomplete category, 3
- codegeneracy operator, 33
- coface operator, 33
- cofibrant
 - object, 46
 - replacement, 49
- cofibration, 46
 - of simplicial sets, 39
- cofinal
 - subset of a poset, 8
- compact object, 11
- compactly-generated Hausdorff space, 38
- complete category, 3
- connected components, 37
- cosimplicial identities, 34
- cylinder object, 49
- degeneracy operator, 35
- directed preorder, 10
- exponential object, *see* cartesian closed category

- face operator, 35
- factorisation system
 - functorial —, 63
 - orthogonal —, 59, 63
 - weak —, 59
- fibrant
 - object, 46
 - replacement, 49
- fibration, 46
 - of simplicial sets, *see* Kan fibration
- filtered category, 10
- fundamental category, 37
- geometric realisation
 - of a simplicial set, 38
- homotopical category, 41
 - saturated —, 41
 - slice —, 41
- homotopical equivalence, 42
- homotopical functor, 41
- homotopical Kan extension, 44
- homotopically contractible, 43
- homotopically initial, 42
- homotopically terminal, 42
- homotopy
 - left —, 50
 - right —, 50
- homotopy category, 54, 65
 - of a model category, 50, 54
- homotopy inverse, 53
- horn, 39
- ind-completion, 13
- ind-object, 13
- Kan complex, 39
- Kan extension, 70
 - absolute —, 70
 - pointwise —, 70
- Kan fibration, 39
 - trivial —, 40
- Kelley space, *see* compactly-generated Hausdorff space
- lifting property, 57
- localisation, 65
- locally presentable category, 16
 - classification theorem, 17
- mate, 24
- model category, 45
 - opposite —, 46
 - slice —, 47
- nerve
 - of a category, 37
- ordinal, 4
- orthogonality, 57
- path object, 49
- pre-universe, 1
- Quillen adjunction, 47, 48
- Quillen equivalence, 47
- Quillen functor, 47
- rank
 - of a set, 6
- relative category, 64
 - maximal —, 64
 - minimal —, 64
 - opposite —, 64
 - saturated —, 41, 65
 - semi-saturated —, 65
- relative functor, 64
- set, 2
- simplex category, 33

- simplicial identities, 35
- simplicial object, 33
- simplicial set, 35
 - discrete —, 37
- singular set, 38
- standard simplex
 - as a simplicial set, 36
 - as a topological space, 37
- transitive set, 5
- trivial cofibration
 - of simplicial sets, *see* anodyne extension
- universe, 1
- universe convention, 4
- weak equivalence, 46, 69
 - natural, 41
 - Quillen —, 40
- well-ordered set
 - classification theorem, 5
- zigzag, 68